arXiv:1211.6732v2 [math.GT] 29 Nov 2012

EQUIVARIANT KHOVANOV-ROZANSKY HOMOLOGY AND LEE-GORNIK SPECTRAL SEQUENCE

HAO WU

Abstract. Lobb observed in [8] that each equivariant $\mathfrak{sl}(N)$ Khovanov-Rozansky homology over $\mathbb{C}[a]$ admits a standard decomposition of a simple form. In the present paper, we derive a formula for the corresponding Lee-Gornik spectral sequence in terms of this decomposition. Moreover, for each equivariant $\mathfrak{sl}(N)$ Khovanov-Rozansky homology over $\mathbb{C}[a]$, we define its torsion width, which provides a lower bound for the homological thickness of the $\mathfrak{sl}(N)$ Khovanov-Rozansky homology and determines when the corresponding Lee-Gornik spectral sequence collapses.

1. Introduction

Our goal is to use the equivariant $\mathfrak{sl}(N)$ Khovanov-Rozansky homology introduced by Krasner in [5] to understand other versions of the $\mathfrak{sl}(N)$ Khovanov-Rozansky homology. Since the algebra is much easier over a principal ideal domain, we work with equivariant $\mathfrak{sl}(N)$ Khovanov-Rozansky homologies over $\mathbb{C}[a]$. Lobb observed in [8] that each such homology admits a standard decomposition of a simple form. The main result of the present paper is a formula for the corresponding Lee-Gornik spectral sequence in terms of this decomposition. Moreover, for each equivariant $\mathfrak{sl}(N)$ Khovanov-Rozansky homology over $\mathbb{C}[a]$, we define a numerical invariant, the torsion width, which provides a lower bound for the homological thickness of the $\mathfrak{sl}(N)$ Khovanov-Rozansky homology and determines exactly when the corresponding Lee-Gornik spectral sequence collapses.

In the remainder of this section, we briefly review the background of this work and state our results.

1.1. **Equivariant $\mathfrak{sl}(N)$ Khovanov-Rozansky homology over $\mathbb{C}[a]$.** Following the grading convention in [4], let x be a homogeneous variable of degree 2 and a a homogeneous variable of degree $2k$, where k is a positive integer. We consider the following homogeneous element of degree $2(N+1)$ of $\mathbb{C}[x,a]$.

$$P(x,a) = x^{N+1} + axF(x,a), \tag{1.1}$$

where $F(x,a)$ is a homogeneous element of $\mathbb{C}[x,a]$ of degree $2(N-k)$.

For any oriented link diagram D, one can use $P(x,a)$ to specialize Krasner's construction in [5] to give a bounded chain complex $C_P(D)$ of graded matrix factorizations over $\mathbb{C}[a]$. We will review the construction of $C_P(D)$ in more details in Section 2. For now, recall that $C_P(D)$ comes with:

- Two $\mathbb{Z}$-gradings: the homological grading and the polynomial grading;
- A filtration: the x-filtration $\mathcal{F}_x$;
- Two differential maps: d_{mf} from the underlying matrix factorizations and d_χ from crossing information.

The homology $H(C_P(D), d_{mf})$ is a finitely generated free $\mathbb{C}[a]$-module that inherits both $\mathbb{Z}$-gradings and the x-filtration. The equivariant $\mathfrak{sl}(N)$ Khovanov-Rozansky homology of D over $\mathbb{C}[a]$ with potential $P(x,a)$ is defined to be the homology

$$H_P(D) = H(H(C_P(D), d_{mf}), d_\chi), \tag{1.2}$$

which, again, inherits both $\mathbb{Z}$-gradings and the x-filtration.

As a special case of Krasner's work in [5], we have the following theorem.

2010 *Mathematics Subject Classification.* Primary 57M27.

Key words and phrases. Equivariant Khovanov-Rozansky homology, homological thickness, Lee-Gornik spectral sequence.

The author was partially supported by NSF grant DMS-1205879 and a Collaboration Grant for Mathematicians from the Simons Foundation.

Theorem 1.1. [5] *Every Reidemeister move of D induces a homotopy equivalence of $C_P(D)$ that preserves both $\mathbb{Z}$-gradings and the x-filtration. Consequently, $H_P(D)$, with its two $\mathbb{Z}$-gradings and x-filtration, is invariant under Reidemeister moves.*

Remark 1.2. Strictly speaking, Krasner only proved the invariance under braid-like Reidemeister moves. But the proof of the invariance under Reidemeister move II_b is very similar and given in [13, Theorem 8.2].

Also, the x-filtration is not mentioned in [5]. But, from its definition in Section 2 below, one can see that the homotopy equivalence associated to Reidemeister moves given in [5, 13] preserve the x-filtration.

Define $C_N(D) = C_P(D)/aC_P(D)$. Then $C_N(D)$ is isomorphic to the $\mathfrak{sl}(N)$ Khovanov-Rozansky chain complex in [4]. It inherits from $C_P(D)$:

- the homological grading and the polynomial grading[1],
- both differential maps: d_{mf} and d_χ.

The homology

$$H_N(D) = H(H(C_N(D), d_{mf}), d_\chi), \tag{1.3}$$

is the $\mathfrak{sl}(N)$ Khovanov-Rozansky homology defined in [4]. The invariance of $H_N(D)$ was established by Khovanov and Rozansky in [4] but can now be viewed as a corollary of Theorem 1.1.

Corollary 1.3. [4] *Every Reidemeister move of D induces a homotopy equivalence of $C_N(D)$ that preserves both the homological grading and the polynomial grading. Consequently, $H_N(D)$, with its homological grading and polynomial grading, is invariant under Reidemeister moves.*

Proof. The standard quotient map $C_P(D) \to C_N(D)$ preserves homotopy equivalence. So Corollary 1.3 follows from Theorem 1.1. $\square$

Define $\hat{C}_P(D) = C_P(D)/(a-1)C_P(D)$. Then $\hat{C}_P(D)$ is a bounded chain complex of filtered matrix factorizations over $\mathbb{C}$. It inherits from $C_P(D)$:

- the homological grading,
- the x-filtration $\mathcal{F}_x$,
- both differential maps: d_{mf} and d_χ.

We call the homology

$$\hat{H}_P(D) = H(H(\hat{C}_P(D), d_{mf}), d_\chi), \tag{1.4}$$

the deformed $\mathfrak{sl}(N)$ Khovanov-Rozansky homology with potential $P(x,1)$. This version of the Khovanov-Rozansky homology was originally introduced by Lee [7] in the $\mathfrak{sl}(2)$ case and then by Gornik [2] in the general $\mathfrak{sl}(N)$ case. Its invariance was first established by the author in [11] but can now be viewed as a corollary of Theorem 1.1.

Corollary 1.4. [11] *Every Reidemeister move of D induces a homotopy equivalence of $\hat{C}_P(D)$ that preserves both the homological grading and the x-filtration. Consequently, $\hat{H}_P(D)$, with its homological grading and x-filtration, is invariant under Reidemeister moves.*

Proof. The standard quotient map $C_P(D) \to \hat{C}_P(D)$ preserves homotopy equivalence. So Corollary 1.4 follows from Theorem 1.1. $\square$

1.2. **The Lee-Gornik spectral sequences.**

Theorem 1.5. [2, 7] *Let D be a diagram of an oriented link L. Then*

- *The x-filtration $\mathcal{F}_x$ on the chain complex $(H(\hat{C}_P(D), d_{mf}), d_\chi)$ induces a spectral sequence $\{\hat{E}_r(L)\}$ converging to $\hat{H}_P(L)$ with $\hat{E}_1(L) \cong H_N(L)$,*
- *The x-filtration $\mathcal{F}_x$ on the chain complex $(H(C_P(D), d_{mf}), d_\chi)$ induces a spectral sequence $\{E_r(L)\}$ converging to $H_P(L)$ with $E_1(L) \cong H_N(L) \otimes_{\mathbb{C}} \mathbb{C}[a]$.*

---

[1] The increasing filtration induced by this polynomial grading is the same as the x-filtration that $C_N(D)$ inherits from $C_P(D)$.

Remark 1.6. Only the E_0-pages of $\{\hat{E}_r(L)\}$ and $\{E_r(L)\}$ depend on the choice of the diagram D. By Theorem 1.1 and Corollary 1.4, for $r \geq 1$, $\hat{E}_r(L)$ and $E_r(L)$ are link invariants.

The spectral sequence $\{\hat{E}_r(L)\}$ was first observed by Lee [7] in the $\mathfrak{sl}(2)$ case and then generalized to the $\mathfrak{sl}(N)$ case by Gornik [2]. A complete construction of $\{\hat{E}_r(L)\}$ can be found in [11]. The construction of $\{E_r(L)\}$ is very similar and given in Section 3 below. We call $\{\hat{E}_r(L)\}$ the Lee-Gornik spectral sequence over $\mathbb{C}$ and $\{E_r(L)\}$ the Lee-Gornik spectral sequence over $\mathbb{C}[a]$.

1.3. **Lobb's decomposition theorem.** As shown in Section 3 below, $(H(C_P(D), d_{mf}), d_\chi)$ is a bounded chain complex of finitely generated graded free $\mathbb{C}[a]$-module. Lobb [8] observed that this implies $(H(C_P(D), d_{mf}), d_\chi)$ decomposes into a direct sum of simple graded chain complexes of the forms

$$F_{i,s} = 0 \to \mathbb{C}[a]\|i\|\{s\} \to 0, \tag{1.5}$$

$$T_{i,m,s} = 0 \to \mathbb{C}[a]\|i-1\|\{s+2km\} \xrightarrow{a^m} \mathbb{C}[a]\|i\|\{s\} \to 0, \tag{1.6}$$

where $\|i\|$ indicates that the component is of homological degree i and, following [4], $\{s\}$ means shifting the polynomial grading up by s. Therefore, $H_P(D)$ is the direct sum of a free graded $\mathbb{C}[a]$-module and torsion components of the form $\mathbb{C}[a]/(a^m)$. The torsion part of $H_P(D)$ is not yet well understood. But the free part of $H_P(D)$ is relatively simple and can be explicitly described using the deformed $\mathfrak{sl}(N)$ Khovanov-Rozansky homology. Theorem 1.7 below is a more precise formulation of the decomposition of $H_P(L)$ observed by Lobb in [8].

For any oriented link L, denote by $\hat{H}_P^i(L)$ the component of $\hat{H}_P(L)$ of homological grading i and by $\hat{\mathcal{H}}_P^i(L)$ the graded $\mathbb{C}$-linear space associated to the filtered space $(\hat{H}_P^i(L), \mathcal{F}_x)$. That is, $\hat{\mathcal{H}}_P^i(L) = \bigoplus_{j\in\mathbb{Z}} \hat{\mathcal{H}}_P^{i,j}(L)$, where $\hat{\mathcal{H}}_P^{i,j}(L) = \mathcal{F}_x^j \hat{H}_P^i(L)/\mathcal{F}_x^{j-1}\hat{H}_P^i(L)$.

Theorem 1.7. [8] *Given an oriented link L and a homological degree i, there is a finite sequence*

$$\{(m_{i,1}, s_{i,1}), \dots, (m_{i,n_i}, s_{i,n_i})\} \subset \mathbb{Z}_{>0} \times \mathbb{Z}$$

such that, as graded $\mathbb{C}[a]$*-modules,*

$$H_P^i(L) \cong (\hat{\mathcal{H}}_P^i(L) \otimes_{\mathbb{C}} \mathbb{C}[a]) \oplus \bigoplus_{l=1}^{n_i} (\mathbb{C}[a]/(a^{m_{i,l}}))\{s_{i,l}\}, \tag{1.7}$$

where $H_P^i(L)$ *is component of* $H_P(L)$ *of homological grading* i*. Moreover, the sequence*

$$\{(m_{i,1}, s_{i,1}), \dots, (m_{i,n_i}, s_{i,n_i})\}$$

is unique up to permutation.

A complete proof of Theorem 1.7 is given in Subsection 4.2 below. A byproduct of this theorem is a decomposition of $H_N(L)$, which we formulate in the following corollary. See Subsection 4.2 below for its proof.

Corollary 1.8. *Using notations in Theorem 1.7, we have*

$$H_N^i(L) \cong \hat{\mathcal{H}}_P^i(L) \oplus \left(\bigoplus_{l=1}^{n_i} \mathbb{C}\{s_{i,l}\}\right) \oplus \left(\bigoplus_{l=1}^{n_{i+1}} \mathbb{C}\{2km_{i+1,l} + s_{i+1,l}\}\right), \tag{1.8}$$

where $H_N^i(L)$ *is component of* $H_N(L)$ *of homological grading* i*.*

1.4. **Decompositions of the Lee-Gornik spectral sequences.** The main results of the present paper are formulas for $\{E_r(L)\}$ and $\{\hat{E}_r(L)\}$ in terms of decomposition (1.7). To state our result, we need to introduce a non-standard tensor product "$\boxtimes$" of bigraded vector spaces.[2]

---

[2] The definition of "$\boxtimes$" in Definition 1.9 comes from the normalization we use in the definition of the spectral sequence of a filtered chain complex. If one uses a different normalization, then the definition of "$\boxtimes$" needs to change accordingly.

Definition 1.9. Let $\mathcal{H} = \bigoplus_{i,j} \mathcal{H}^{i,j}$ and $E = \bigoplus_{p,q} E^{p,q}$ be two $\mathbb{Z}^{\oplus 2}$-graded $\mathbb{C}$-spaces. Then

$$\mathcal{H} \boxtimes E = \bigoplus_{\alpha,\beta} (\mathcal{H} \boxtimes E)^{\alpha,\beta}$$

is the $\mathbb{Z}^{\oplus 2}$-graded $\mathbb{C}$-space satisfying

$$(\mathcal{H} \boxtimes E)^{\alpha,\beta} = \bigoplus_{j+p=\alpha,\ q+i-j=\beta} \mathcal{H}^{i,j} \otimes_{\mathbb{C}} E^{p,q}.$$

Next, we define the x-filtration $\mathcal{F}_x$ of $F_{i,s}$, $T_{i,m,s}$, $\hat{F}_{i,s} = F_{i,s}/(a-1)F_{i,s}$ and $\hat{T}_{i,m,s} = T_{i,m,s}/(a-1)T_{i,m,s}$.

$$\mathcal{F}_x^p F_{i,s} = \begin{cases} 0 \to \mathbb{C}[a]\|i\|\{s\} \to 0 & \text{if } p \geq s, \\ 0 & \text{if } p < s. \end{cases} \tag{1.9}$$

$$\mathcal{F}_x^p T_{i,m,s} = \begin{cases} 0 \to \mathbb{C}[a]\|i-1\|\{s+2km\} \xrightarrow{a^m} \mathbb{C}[a]\|i\|\{s\} \to 0 & \text{if } p \geq s+2km \\ 0 \to \mathbb{C}[a]\|i\|\{s\} \to 0 & \text{if } s \leq p < s+2km \\ 0 & \text{if } p < s. \end{cases} \tag{1.10}$$

$$\mathcal{F}_x^p \hat{F}_{i,s} = \begin{cases} 0 \to \mathbb{C}\|i\| \to 0 & \text{if } p \geq s, \\ 0 & \text{if } p < s, \end{cases} \tag{1.11}$$

$$\mathcal{F}_x^p \hat{T}_{i,m,s} = \begin{cases} 0 \to \mathbb{C}\|i-1\| \xrightarrow{1} \mathbb{C}\|i\| \to 0 & \text{if } p \geq s+2km \\ 0 \to \mathbb{C}\|i\| \to 0 & \text{if } s \leq p < s+2km \\ 0 & \text{if } p < s, \end{cases} \tag{1.12}$$

The filtered chain complexes $F_{i,s}$, $T_{i,m,s}$, $\hat{F}_{i,s}$ and $\hat{T}_{i,m,s}$ are very simple. Their spectral sequences are given in the following lemma, which is proved in Subsection 4.3 below.

Lemma 1.10. *For any $r \geq 0$,*

$$E_r^{p,q}(F_{i,s}) \cong \begin{cases} \mathbb{C}[a]\{s\} & \text{if } p = s \text{ and } q = i-s, \\ 0 & \text{otherwise,} \end{cases} \tag{1.13}$$

$$E_r^{p,q}(T_{i,m,s}) \cong \begin{cases} (\mathbb{C}[a]/(a^m))\{s\} & \text{if } p = s,\ q = i-s \text{ and } r \geq 2km+1, \\ \mathbb{C}[a]\{s\} & \text{if } p = s,\ q = i-s \text{ and } r \leq 2km, \\ \mathbb{C}[a]\{s+2km\} & \text{if } p = s+2km,\ q = i-1-s-2km \text{ and } r \leq 2km, \\ 0 & \text{otherwise,} \end{cases} \tag{1.14}$$

$$E_r^{p,q}(\hat{F}_{i,s}) \cong \begin{cases} \mathbb{C} & \text{if } p = s \text{ and } q = i-s, \\ 0 & \text{otherwise.} \end{cases} \tag{1.15}$$

$$E_r^{p,q}(\hat{T}_{i,m,s}) \cong \begin{cases} \mathbb{C} & \text{if } p = s,\ q = i-s \text{ and } r \leq 2km, \\ \mathbb{C} & \text{if } p = s+2km,\ q = i-1-s-2km \text{ and } r \leq 2km, \\ 0 & \text{otherwise.} \end{cases} \tag{1.16}$$

Note that:

- *Isomorphisms (1.13) and (1.14) preserve the polynomial grading.*
- *Both $\{E_r(T_{i,m,s})\}$ and $\{E_r(\hat{T}_{i,m,s})\}$ collapse exactly at their E_{2km+1}-pages.*[3]
- *Both $\{E_r(F_{i,s})\}$ and $\{E_r(\hat{F}_{i,s})\}$ collapse at their E_0-pages.*
- *$E_r(F_{i,s}) \cong \mathbb{C}\|i\|\{s\} \boxtimes E_r(F_{0,0})$, $E_r(\hat{F}_{i,s}) \cong \mathbb{C}\|i\|\{s\} \boxtimes E_r(\hat{F}_{0,0})$, where "$\boxtimes$" is the product defined in Definition 1.9 and $\mathbb{C}\|i\|\{s\}$ is the $\mathbb{Z}^{\oplus 2}$-graded $\mathbb{C}$-space given by*

$$(\mathbb{C}\|i\|\{s\})^{p,q} = \begin{cases} \mathbb{C} & \text{if } p = i \text{ and } q = s, \\ 0 & \text{otherwise.} \end{cases}$$

---

[3] We say that a spectral sequence $\{E_r\}$ collapses exactly at its E_t-page if $E_{t-1} \ncong E_t$ but $E_{t+r} \cong E_t$ $\forall r \geq 0$.

Combining Lemma 1.10 and the following theorem, we get explicit formulas for $\{E_r(L)\}$ and $\{\hat{E}_r(L)\}$ in terms of Lobb's decomposition (Theorem 1.7.)

Theorem 1.11. *For an oriented link L, let $\hat{\mathcal{H}}_P(L) = \bigoplus_{i\in\mathbb{Z}} \hat{\mathcal{H}}_P^i(L) = \bigoplus_{(i,j)\in\mathbb{Z}^{\oplus 2}} \hat{\mathcal{H}}_P^{i,j}(L)$ and, for each i,*

$$\{(m_{i,1}, s_{i,1}), \dots, (m_{i,n_i}, s_{i,n_i})\} \subset \mathbb{Z}_{>0} \times \mathbb{Z}$$

the sequence given in Theorem 1.7. Then, for any $r \geq 1$,

$$\hat{E}_r(L) \quad\cong\quad (\hat{\mathcal{H}}_P(L) \boxtimes E_r(\hat{F}_{0,0})) \oplus \bigoplus_{i\in\mathbb{Z}} \bigoplus_{l=1}^{n_i} E_r(\hat{T}_{i,m_{i,l},s_{i,l}}), \tag{1.17}$$

$$E_r(L) \quad\cong\quad (\hat{\mathcal{H}}_P(L) \boxtimes E_r(F_{0,0})) \oplus \bigoplus_{i\in\mathbb{Z}} \bigoplus_{l=1}^{n_i} E_r(T_{i,m_{i,l},s_{i,l}}), \tag{1.18}$$

where isomorphism (1.17) *preserves the usually (p,q)-grading of spectral sequences, while isomorphism* (1.18) *preserves the usually (p,q)-grading of spectral sequences as well as the polynomial grading of each $E_r^{p,q}$-component.*

Theorem 1.11 is proved in Subsection 4.3 below. The key to its proof is that, when the chain complex $(H(C_P(D), d_{mf}), d_\chi)$ is decomposed into complexes of the forms $F_{i,s}$ and $T_{i,m,s}$, the x-filtration decomposes accordingly. This is established in Subsection 4.1.

Remark 1.12. With minor cosmetic changes, all Theorems 1.7, 1.11, Corollary 1.8 and their proofs generalize to the colored $\mathfrak{sl}(N)$ link homology defined in [12, 13, 14].

1.5. The torsion width of $H_P(L)$.

Definition 1.13. Let L be an oriented link. Using the notations in Theorem 1.7, we define the torsion width of $H_P(L)$ to be $\mathrm{tw}_P(L) = \max\{2km_{i,l} \mid i \in \mathbb{Z}, 1 \leq l \leq n_i\}$[4], which, by Theorem 1.7, is a link invariant. Equivalently, one has $\mathrm{tw}_P(L) = \min\{2km \mid m \in \mathbb{Z}_{\geq 0},\ a^m H_P(L) \text{ is free}\}$.

Here, recall that $2k$ is the degree of a. We include the factor $2k$ in the definition of $\mathrm{tw}_P(L)$ to make it invariant under simple changes of variable of the form $a = b^j$.

Corollary 1.14. *Let L be an oriented link with torsion width $\mathrm{tw}_P(L) = w$, and D a diagram of L. Assume D is not a union of disjoint circles embedded in the plane. Then both spectral sequences $\{E_r(L)\}$ and $\{\hat{E}_r(L)\}$ from Theorem 1.5 collapse exactly at their E_{w+1}-pages.*

Proof. From Lemma 1.10, we know that $\{E_r(F_{i,s})\}$ and $\{E_r(\hat{F}_{i,s})\}$ both collapse exactly at their E_0-pages, while $\{E_r(T_{i,m,s})\}$ and $\{E_r(\hat{T}_{i,m,s})\}$ both collapse exactly at their E_{2km+1}-pages. So this corollary follows from Theorem 1.11. □

Definition 1.15. For an oriented link L, denote by $H_N^{i,j}(L)$ the component of $H_N(L)$ of homological degree i and polynomial degree j. Define the $\mathfrak{sl}(N)$ homological thickness $\mathrm{ht}_N(L)$ and the local $\mathfrak{sl}(N)$ homological thickness $\mathrm{lht}_N(L)$ of L to be

$$\mathrm{ht}_N(L) \quad=\quad \max\{1 + \frac{1}{2}[(2i_1 + j_1) - (2i_2 + j_2)] \mid H_N^{i_1,j_1}(L) \neq 0,\ H_N^{i_2,j_2}(L) \neq 0,\} \tag{1.19}$$

$$\mathrm{lht}_N(L) \quad=\quad \max\{\frac{j_1 - j_2}{2} \mid \exists i \in \mathbb{Z},\ \text{such that } H_N^{i,j_1}(L) \neq 0,\ H_N^{i+1,j_2}(L) \neq 0.\} \tag{1.20}$$

Of course, $\mathrm{ht}_N(L)$ is a naive generalization of the homological thickness of the rational Khovanov homology. From their definitions, one can see that $\mathrm{lht}_N(L) \leq \mathrm{ht}_N(L)$ for any L. See [1, Table 2] for an example of $\mathrm{lht}_2(L) < \mathrm{ht}_2(L)$.

Corollary 1.16. *If $H_N(L) \ncong \hat{\mathcal{H}}_P(L) := \bigoplus_{i\in\mathbb{Z}} \hat{\mathcal{H}}_P^i(L)$, then $\mathrm{tw}_P(L) \leq 2\mathrm{lht}_N(L) \leq 2\mathrm{ht}_N(L)$.*

Proof. By Corollary 1.8, each torsion component $(\mathbb{C}[a]/(a^m))\|i\|\{s\}$ of $H_P(L)$ generates a pair of 1-dimensional components of $H_N(L)$: $\mathbb{C}\|i\|\{s\}$ and $\mathbb{C}\|i-1\|\{2km+s\}$. Corollary 1.16 follows from this observation. □

---

[4] We use the convention that $\mathrm{tw}_P(L) = 0$ if $H_P(L)$ is a free $\mathbb{C}[a]$-module.

The pairing of $\mathbb{C}\|i\|\{s\}$ and $\mathbb{C}\|i-1\|\{2km+s\}$ is a generalization of [7, Theorem 1.4], which states that, except those with homological degree 0, all homogeneous generators of the rational Khovanov homology of an alternating knot are paired into pairs of bi-degree difference $(-1,4)$. In fact, we can use the torsion width to recover this theorem. We state this in the following corollary.

Corollary 1.17. [7, Theorem 1.4] *Suppose* $N=2$, $\deg a = 4$ *and* $P(x,a)=x^3-ax$. *Assume the knot* K *is not the unknot and* $\mathrm{lht}_2(K)=2$. *Then* $\mathrm{tw}_P(K)=4$ *and, consequently, there exist pairs of integers* $(i_1,s_1),\dots,(i_n,s_n)$ *such that* $H_2(K)\cong\hat{\mathcal{H}}_P(K)\oplus\bigoplus_{l=1}^{n}(\mathbb{C}\|i_l\|\{s_l\}\oplus\mathbb{C}\|i_l-1\|\{s_l+4\})$.

Proof. If $H_P(K)$ is a free $\mathbb{C}[a]$-module, then, by Corollary 1.8, $H_2(K)\cong\hat{\mathcal{H}}_P(K)$, which is 2-dimensional by [7, Theorem 4.2]. Thus, by [6, Theorem 1.1], K is the unknot, which contradicts our assumption. So $H_P(K)$ has torsion. Since $\deg a=4$, its torsion width is at least 4. On the other hand, by Corollary 1.16, we have $\mathrm{tw}_P(K)\leq 2\mathrm{lht}_2(K)=4$. Therefore, $\mathrm{tw}_P(K)=4$, which means all torsion components of $H_P(K)$ are of the form $\mathbb{C}[a]/(a)\|i\|\{s\}$. The decomposition of $H_2(K)$ in the corollary then follows from Corollary 1.8. □

1.6. **Organization of this paper.** We review the constructions of H_P, $E_r(L)$ and $\hat{E}_r(L)$ in Sections 2 and 3. Then we prove the decomposition theorems in Section 4.

We assume the reader is somewhat familiar with the construction of the $\mathfrak{sl}(N)$ Khovanov-Rozansky homology in [4].

2. Definition of H_P

In the remainder of this paper, N and k are fixed positive integers with $N\geq k$.

2.1. **Graded and filtered matrix factorizations.** Let $R=\mathbb{C}[x_1,\dots,x_m,a]$, where $x_1,\dots,x_m$ are homogeneous variables of degree 2 and a is a homogeneous variable of degree $2k$. We endow two structures on R:

- The polynomial grading with degree function deg given by $\deg(a^l\prod_{i=1}^m x_i^{l_i})=2kl+\sum_{i=1}^m 2l_i$.
- The x-filtration $0=\mathcal{F}_x^{-1}R\subset\mathcal{F}_x^0R\subset\cdots\subset\mathcal{F}_x^nR\subset\cdots$ such that $a^l\prod_{i=1}^m x_i^{l_i}\in\mathcal{F}_x^nR$ if and only if $\sum_{i=1}^m 2l_i\leq n$. The degree function $\deg_x$ of $\mathcal{F}_x$ is given by $\deg_x(a^l\prod_{i=1}^m x_i^{l_i})=\sum_{i=1}^m 2l_i$.

Unless otherwise specified, when we say an element is homogeneous, we mean it is homogeneous with respect to the polynomial grading.

Definition 2.1. Let M be an R-module. We say that M is a graded R-module if it is endowed with a grading $M=\bigoplus_i M_i$ such that, for any homogeneous element r of R, $rM_i\subset M_{i+\deg r}$. We say that M is an x-filtered R-module if it is endowed with an increasing filtration $\mathcal{F}_x$ such that, for any element r of R, $r\mathcal{F}_x^iM\subset\mathcal{F}_x^{i+\deg_x r}M$.

Definition 2.2. Let w be a homogeneous element of R with $\deg w=2N+2$. A matrix factorization M of w over R is a collection of two free R-modules M^0, M^1 and two R-module homomorphisms $d^0:M^0\to M^1$, $d^1:M^1\to M^0$, called differential maps, such that $d^1d^0=w\cdot\mathrm{id}_{M^0}$ and $d^0d^1=w\cdot\mathrm{id}_{M^1}$. We usually write M as $M^0\xrightarrow{d^0}M^1\xrightarrow{d^1}M^0$.

We call M graded if M^0, M^1 are graded R-modules and d^0, d^1 are homogeneous homomorphisms with $\deg d^0=\deg d^1=N+1$.

We call M x-filtered if M^0 and M^1 are x-filtered R-modules and $\deg_x d^0,\deg_x d^1\leq N+1$.

In the definition of H_P, we use only Koszul matrix factorizations defined below.

Definition 2.3. Let b and c be homogeneous elements of R with $\deg(bc)=2N+2$. Denote by $(b,c)_R$ the Koszul matrix factorization

$$R\xrightarrow{b}R\{N+1-\deg b\}\xrightarrow{c}R,$$

where b, c act on R by multiplication and "$\{s\}$" means shifting by s both the polynomial grading and the x-filtration[5] of R. This matrix factorization of bc is both graded and x-filtered.

---

[5] That is, in $R\{s\}$, $\deg(a^l\prod_{i=1}^m x_i^{l_i})=s+2kl+\sum_{i=1}^m 2l_i$ and $\deg_x(a^l\prod_{i=1}^m x_i^{l_i})=s+\sum_{i=1}^m 2l_i$.

For homogeneous elements $b_1 \cdots, b_n, c_1, \cdots, c_n$ of R with $\deg(b_i c_i) = 2N + 2$, $i = 1, \cdots, n$, denote by

$$\left(\begin{array}{cc} b_1 & c_1 \\ b_2 & c_2 \\ \vdots & \vdots \\ b_n & c_n \end{array}\right)_R$$

the Koszul matrix factorization $(b_1, c_1)_R \otimes_R (b_2, c_2)_R \otimes_R \cdots \otimes_R (b_n, c_n)_R$. This matrix factorization of $w = \sum_{i=1}^n b_i c_i$ is again both graded and x-filtered.

When R is clear from context, we drop it from the notation.

Definition 2.4. As a free R-module, $(b, c)_R$ has a basis $\{1_0, 1_1\}$, where 1_ε is the "1" in the copy of R with $\mathbb{Z}_2$-grading ε. More generally, the tensor product

$$\left(\begin{array}{cc} b_1 & c_1 \\ b_2 & c_2 \\ \vdots & \vdots \\ b_n & c_n \end{array}\right)_R = (b_1, c_1)_R \otimes_R (b_2, c_2)_R \otimes_R \cdots \otimes_R (b_n, c_n)_R$$

has a basis $\{1_{\vec{\varepsilon}} \mid \vec{\varepsilon} = (\varepsilon_1, \ldots, \varepsilon_n) \in \mathbb{Z}_2^n\}$, where $1_{\vec{\varepsilon}} = 1_{\varepsilon_1} \otimes \cdots \otimes 1_{\varepsilon_n}$. We call $\{1_{\vec{\varepsilon}}\}$ the standard basis for this Koszul matrix factorization.

Note that

- $\{1_{\vec{\varepsilon}}\}$ is a homogeneous basis with respect to the polynomial grading.
- $\deg 1_{\vec{\varepsilon}} = \deg_x 1_{\vec{\varepsilon}}$ for every $\vec{\varepsilon}$.
- $\deg_x(\sum_{\vec{\varepsilon}} f_{\vec{\varepsilon}} 1_{\vec{\varepsilon}}) \leq l$ if and only if $\deg_x f_{\vec{\varepsilon}} \leq l - \deg_x 1_{\vec{\varepsilon}}$ for every $\vec{\varepsilon}$.

The following simple observation is very useful in the proof of the decomposition theorems in Section 4.

Lemma 2.5. *Suppose M is a Koszul matrix factorization as defined above and ρ is a homogeneous element of M. Then $\deg_x \rho \leq \deg \rho$, and $\deg_x \rho < \deg \rho$ if and only if $\rho \in aM$.*

Proof. Let $\{1_{\vec{\varepsilon}}\}$ be the standard basis for M. Then $\rho = \sum_{\vec{\varepsilon}} f_{\vec{\varepsilon}} 1_{\vec{\varepsilon}}$, where $f_{\vec{\varepsilon}}$ is a homogeneous element of R with $\deg f_{\vec{\varepsilon}} = \deg \rho - \deg 1_{\vec{\varepsilon}}$. Note that, for every $f \in R$, $\deg_x f \leq \deg f$, and $\deg_x f < \deg f$ if and only if $f \in aR$. So $\deg_x \rho = \max_{\vec{\varepsilon}}(\deg_x f_{\vec{\varepsilon}} + \deg_x 1_{\vec{\varepsilon}}) \leq \max_{\vec{\varepsilon}}(\deg f_{\vec{\varepsilon}} + \deg 1_{\vec{\varepsilon}}) = \deg \rho$. Moreover, $\deg_x \rho < \deg \rho$ if and only if $\deg_x f_{\vec{\varepsilon}} < \deg f_{\vec{\varepsilon}}$ for every $\vec{\varepsilon}$ if and only if $f_{\vec{\varepsilon}} \in aR$ for every $\vec{\varepsilon}$. □

2.2. The matrix factorization associated to a MOY graph.

Definition 2.6. A MOY graph Γ is a finite oriented graph embedded in $\mathbb{R}^2$ with the following properties:

1. Edges of Γ are divided into two types: regular edges and wide edges.
2. Vertices of Γ are of two types:
    - endpoints: 1-valent vertices that are endpoints of regular edges,
    - internal vertices: 3-valent vertices with
        - either two regular edges pointing inward and one wide edge pointing outward,
        - or two regular edges pointing outward and one wide edge pointing inward.

We say that Γ is closed if it has no endpoints.

A marking of Γ consists of

1. A finite set of marked points on Γ such that:
    - every regular edge contains at least one marked point, no wide edges contain any marked points,
    - every endpoint of Γ is marked, none of the internal vertices are marked.
2. An assignment that assigns to each marked point a different homogeneous variable of degree 2.

In the rest of this subsection, we fix a MOY graph Γ and a marking of Γ. Assume $x_1, \ldots, x_m$ are the variables assigned to the marked points of Γ. Define $R = \mathbb{C}[x_1, \ldots, x_m, a]$, where a is a homogeneous variable of degree $2k$. Let $P(x, a)$ be the homogeneous polynomial given in (1.1).

Now cut Γ at all the marked points. This cuts Γ into a collect of simple marked MOY graphs $\Gamma_1, \ldots, \Gamma_n$, each of which is of one of the two types in Figure 1



$\Gamma_{i;p}$ $\Gamma_{i,j;p,q}$

FIGURE 1. Pieces of Γ

Define

$$v_{i;p} = \frac{P(x_i,a) - P(x_p,a)}{x_i - x_p}. \tag{2.1}$$

Since $P(x_i,a) + P(x_j,a)$ is symmetric in x_i and x_j, there is a unique polynomial $G(X,Y,a)$ satisfying $G(x_i+x_j, x_ix_j, a) = P(x_i,a) + P(x_j,a)$. Define

$$u^{\langle 1\rangle}_{i,j;p,q} = \frac{G(x_i+x_j, x_ix_j, a) - G(x_p+x_q, x_ix_j, a)}{x_i+x_j-x_p-x_q}, \tag{2.2}$$

$$u^{\langle 2\rangle}_{i,j;p,q} = \frac{G(x_p+x_q, x_ix_j, a) - G(x_p+x_q, x_px_q, a)}{x_ix_j - x_px_q}. \tag{2.3}$$

Note that $v_{i;j}$, $u^{\langle 1\rangle}_{i,j;p,q}$, $u^{\langle 2\rangle}_{i,j;p,q}$ are all homogeneous elements of R.

Definition 2.7.

$$C_P(\Gamma_{i;p}) := (v_{i;p}, x_i - x_p)_R, \tag{2.4}$$

$$C_P(\Gamma_{i,j;p,q}) := \begin{pmatrix} u^{\langle 1\rangle}_{i,j;p,q} & x_i+x_j-x_p-x_q \\ u^{\langle 2\rangle}_{i,j;p,q} & x_ix_j - x_px_q \end{pmatrix}_R \{-1\}, \tag{2.5}$$

$$C_P(\Gamma) := C_P(\Gamma_1) \otimes_R C_P(\Gamma_2) \otimes_R \cdots \otimes_R C_P(\Gamma_n). \tag{2.6}$$

Note that

1. $C_P(\Gamma_{i;p})$ is a Koszul matrix factorization of $w_{i;p} = P(x_i,a) - P(x_p,a)$.
2. $C_P(\Gamma_{i,j;p,q})$ is a Koszul matrix factorization of $w_{i,j;p,q} = P(x_i,a) + P(x_j,a) - P(x_p,a) - P(x_q,a)$.
3. $C_P(\Gamma)$ is a Koszul matrix factorization of

   $$w = \sum_{x_i \text{ is assigned to an endpoint.}} \pm P(x_i,a),$$

   where the sign is positive if Γ points towards the corresponding endpoint and is negative if Γ points away from the corresponding endpoint. In particular, $w = 0$ if Γ is closed.

Definition 2.8. For a closed MOY graph Γ,

1. define $H_P(\Gamma)$ to be the homology of $C_P(\Gamma)$, which inherits the polynomial grading, the x-filtration and the $\mathbb{Z}_2$-grading of $C_P(\Gamma)$,
2. define $H_N(\Gamma)$ to be the homology of $C_N(\Gamma) := C_P(\Gamma)/aC_P(\Gamma)$, which inherits the polynomial grading and the $\mathbb{Z}_2$-grading of $C_N(\Gamma)$,
3. define $\hat{H}_P(\Gamma)$ to be the homology of $\hat{C}_P(\Gamma) := C_P(\Gamma)/(a-1)C_P(\Gamma)$, which inherits the x-filtration and the $\mathbb{Z}_2$-grading of $C_N(\Gamma)$.

2.3. **The chain complex associated to a link diagram.** A marking of a link diagram D consists of

1. a finite collection of marked points on D such that none of the crossings are marked and every arc between two crossings contains at least one marked point,
2. an assignment that assigns to each marked point a different homogeneous variable of degree 2.

Let D be an oriented link diagram with a marking. Assume $x_1, \dots, x_m$ are the variables assigned to the marked points of D. Define $R = \mathbb{C}[x_1, \dots, x_m, a]$, where a is a homogeneous variable of degree $2k$. Cut D at its marked points. This cuts D into simple pieces $D_1, \dots, D_n$ of the types shown in Figure 2.



FIGURE 2. Pieces of D

We define the chain complex $C_P(\Gamma_{i;p})$ to be

$$C_P(\Gamma_{i;p}) = 0 \to C_P(\Gamma_{i;p})\|0\| \to 0, \tag{2.7}$$

where the term "$C_P(\Gamma_{i;p})$" on the right hand side is the Koszul matrix factorization defined in Definition 2.7 and "$\|0\|$" means this term has homological degree 0.

To define $C_P(c^{\pm}_{i,j;p,q})$, we need the following lemma.



FIGURE 3. Homomorphisms χ_0 and χ_1

Lemma 2.9. [4, 5, 13] *Up to homotopy and scaling, there is a unique homotopically non-trivial homomorphism* $\chi_0 : C_P(\Gamma_{i;p} \sqcup \Gamma_{j;q}) \to C_P(\Gamma_{i,j;p,q})$ *with* $\deg \chi_0 = 1$. *And, up to homotopy and scaling, there is a unique homotopically non-trivial homomorphism* $\chi_1 : C_P(\Gamma_{i,j;p,q}) \to C_P(\Gamma_{i;p} \sqcup \Gamma_{j;q})$ *with* $\deg \chi_1 = 1$. *Moreover, these homomorphisms satisfy* $\deg_x \chi_0 = \deg_x \chi_1 = 1$ *and, up to scaling,* $\chi_1 \circ \chi_0 \simeq (x_p - x_j)\mathrm{id}_{C_P(\Gamma_{i;p} \sqcup \Gamma_{j;q})}$, $\chi_0 \circ \chi_1 \simeq (x_p - x_j)\mathrm{id}_{C_P(\Gamma_{i,j;p,q})}$.

Proof. The uniqueness of χ_0 and χ_1 is proved in a more general setting in [13, Lemma 4.13]. Here, we only recall the explicit construction of χ_0 and χ_1 given by Krasner in [5], which is a straightforward generalization of the corresponding construction by Khovanov and Rozansky in [4].

Recall that

$$C_P(\Gamma_{i;p} \sqcup \Gamma_{j;q}) = \begin{pmatrix} v_{i;p} & x_i - x_p \\ v_{j;q} & x_j - x_q \end{pmatrix}_R = \begin{bmatrix} R \\ R\{2-2n\} \end{bmatrix} \xrightarrow{d^0} \begin{bmatrix} R\{1-n\} \\ R\{1-n\} \end{bmatrix} \xrightarrow{d^1} \begin{bmatrix} R \\ R\{2-2n\} \end{bmatrix},$$

where $d^0 = \begin{pmatrix} v_{i;p} & x_j - x_q \\ v_{j;q} & -x_i + x_p \end{pmatrix}$, $d^1 = \begin{pmatrix} x_i - x_p & x_j - x_q \\ v_{j;q} & -v_{i;p} \end{pmatrix}$, and that

$$C_P(\Gamma_{i,j;p,q}) = \begin{pmatrix} u^{\langle 1\rangle}_{i,j;p,q} & x_i + x_j - x_p - x_q \\ u^{\langle 2\rangle}_{i,j;p,q} & x_i x_j - x_p x_q \end{pmatrix}_R \{-1\} = \begin{bmatrix} R\{-1\} \\ R\{3-2n\} \end{bmatrix} \xrightarrow{\delta^0} \begin{bmatrix} R\{-n\} \\ R\{2-n\} \end{bmatrix} \xrightarrow{\delta^1} \begin{bmatrix} R\{-1\} \\ R\{3-2n\} \end{bmatrix},$$

where $\delta^0 = \begin{pmatrix} u^{\langle 1\rangle}_{i,j;p,q} & x_i x_j - x_p x_q \\ u^{\langle 2\rangle}_{i,j;p,q} & -x_i - x_j + x_p + x_q \end{pmatrix}$,$\delta^1 = \begin{pmatrix} x_i + x_j - x_p - x_q & x_i x_j - x_p x_q \\ u^{\langle 2\rangle}_{i,j;p,q} & -u^{\langle 1\rangle}_{i,j;p,q} \end{pmatrix}$.

In the above explicit forms of $C_P(\Gamma_{i;p} \sqcup \Gamma_{j;q})$ and $C_P(\Gamma_{i,j;p,q})$, define $\chi_0 : C_P(\Gamma_{i;p} \sqcup \Gamma_{j;q}) \to C_P(\Gamma_{i,j;p,q})$ by the matrices $\chi_0^0 = \begin{pmatrix} x_p - x_j & 0 \\ z & 1 \end{pmatrix}$ and $\chi_0^1 = \begin{pmatrix} x_p & -x_j \\ -1 & 1 \end{pmatrix}$, and define $\chi_1 : C_P(\Gamma_{i,j;p,q}) \to C_P(\Gamma_{i;p} \sqcup \Gamma_{j;q})$ by the matrices $\chi_1^0 = \begin{pmatrix} 1 & 0 \\ -z & x_p - x_j \end{pmatrix}$ and $\chi_1^1 = \begin{pmatrix} 1 & x_j \\ 1 & x_p \end{pmatrix}$, where $z = -u^{\langle 2\rangle}_{i,j;p,q} + \frac{u^{\langle 1\rangle}_{i,j;p,q} + x_i u^{\langle 2\rangle}_{i,j;p,q} - v_{j;q}}{x_i - x_p}$. It is straightforward to verify that χ_0 and χ_1 satisfy all the properties in the lemma. □

We define

$$C_P(c^+_{i,j;p,q}) = 0 \to C_P(\Gamma_{i,j;p,q})\|-1\|\{N\} \xrightarrow{\chi_1} C_P(\Gamma_{i;p} \sqcup \Gamma_{j;q})\|0\|\{N-1\} \to 0, \tag{2.8}$$

$$C_P(c^-_{i,j;p,q}) = 0 \to C_P(\Gamma_{i;p} \sqcup \Gamma_{j;q})\|0\|\{1-N\} \xrightarrow{\chi_0} C_P(\Gamma_{i,j;p,q})\|1\|\{-N\} \to 0. \tag{2.9}$$

Definition 2.10. $C_P(D) = C_P(D_1) \otimes_R \cdots \otimes_R C_P(D_n)$, where $C_P(D_i)$ is defined in (2.7), (2.8) and (2.9).

We call the resolution $c^\pm_{i,j;p,q} \rightsquigarrow \Gamma_{i;p} \sqcup \Gamma_{j;q}$ a 0-resolution and $c^\pm_{i,j;p,q} \rightsquigarrow \Gamma_{i,j;p,q}$ a (±1)-resolution. If we choose a 0- or (±1)-resolution for each crossing in D, then we get a MOY graph, which we call a MOY resolution of D. Of course, the marking of D induces a marking of each MOY resolution of D. Let $\mathcal{MOY}(D)$ be the set of all MOY resolutions of D. Denote by w the writhe of D. For each $\Gamma \in \mathcal{MOY}(D)$, let

$$\hbar(\Gamma) = (\#\text{ of }(+1)\text{-resolutions in }\Gamma) - (\#\text{ of }(-1)\text{-resolutions in }\Gamma).$$

Then, as $\mathbb{Z}^{\oplus 2}$-graded R-modules,

$$C_P(D) \cong \bigoplus_{\Gamma \in \mathcal{MOY}(D)} C_P(\Gamma)\|-\hbar(\Gamma)\|\{(N-1)w + \hbar(\Gamma)\}. \tag{2.10}$$

Note that every MOY resolution Γ of D is a closed MOY graph. So $C_P(\Gamma)$ is a Koszul matrix factorization of 0 and, therefore, a $\mathbb{Z}_2$-graded chain complex.[6] Thus, the differential maps of the matrix factorizations of the MOY resolutions of D give rise to a differential map d_{mf} on $C_P(D)$ satisfying:

- d_{mf} is homogeneous with $\deg d_{mf} = \deg_x d_{mf} = N+1$,
- d_{mf} preserves the homological grading.

The differential maps of $C_P(D_i)$ give rise to a differential map d_χ of $C_P(D)$ satisfying:

- d_χ is homogeneous with $\deg d_\chi = \deg_x d_\chi = 0$,
- d_χ raises the homological grading by 1.

As in (1.2), $H_P(D)$ is defined to be $H_P(D) = H(H(C_P(D), d_{mf}), d_\chi)$, which inherits both $\mathbb{Z}$-gradings and the x-filtration of $C_P(D)$. The invariance of $H_P(D)$ is stated in Theorem 1.1.

3. The Lee-Gornik Spectral Sequence

In this section we review the construction of $\hat{E}_r(L)$ and construct $E_r(L)$.

3.1. **Structure of $H_P(\Gamma)$.**

Let Γ be a closed MOY graph with a marking. As before, assume $x_1, \dots, x_m$ are the variables assigned to the marked points of Γ and define $R = \mathbb{C}[x_1, \dots, x_m, a]$, where a is a homogeneous variable of degree $2k$. $P(x,a)$ is the homogeneous polynomial given in (1.1).

If we replace every wide edge in Γ by a pair of parallel regular edges, that is, change $\Gamma_{i,j;p,q}$ to $\Gamma_{i;p} \sqcup \Gamma_{j;q}$ in Figure 3, then we change Γ into a collection of oriented circles embedded in the plane. Denote by ε the total rotation number of this collection and call ε the rotation number of Γ. Furthermore, denote by $H_P^\varepsilon(\Gamma)$ (resp. $\hat{H}_P^\varepsilon(\Gamma)$) the component of $H_P(\Gamma)$ (resp. $\hat{H}_P(\Gamma)$) of $\mathbb{Z}_2$-grading ε and by $H_N^{\varepsilon,n}(\Gamma)$ the component of $H_N(\Gamma)$ of $\mathbb{Z}_2$-grading ε and polynomial grading n.

Lemma 3.1. [2, Proposition 3.2] *As $\mathbb{C}$-linear spaces,*

$$\hat{H}_P^{\varepsilon+1}(\Gamma) \cong 0, \tag{3.1}$$

$$\mathcal{F}_x^n \hat{H}_P^\varepsilon(\Gamma) / \mathcal{F}_x^{n-1} \hat{H}_P^\varepsilon(\Gamma) \cong H_N^{\varepsilon,n}(\Gamma). \tag{3.2}$$

See for example [11, Proposition 2.19] for a complete proof of Lemma 3.1. Slightly modifying this proof, we get Lemma 3.2, which is mentioned in [15] without proof. Since certain technical aspects of its proof are needed later on, we prove Lemma 3.2 in details here.

Lemma 3.2. *As graded $\mathbb{C}[a]$-modules,*

$$H_P^{\varepsilon+1}(\Gamma) \cong 0, \tag{3.3}$$

$$\mathcal{F}_x^n H_P^\varepsilon(\Gamma) / \mathcal{F}_x^{n-1} H_P^\varepsilon(\Gamma) \cong H_N^{\varepsilon,n}(\Gamma) \otimes_{\mathbb{C}} \mathbb{C}[a]. \tag{3.4}$$

---

[6] $C_P(D)$ and $H_P(D)$ both inherit this $\mathbb{Z}_2$-grading. But this $\mathbb{Z}_2$-grading on $H_P(D)$ is always pure and equal to the number of Seifert circles of D. So, unless otherwise specified, we do not keep track of this grading.

Proof. Note that $C_P(\Gamma)$ is also a graded $\mathbb{C}[x_1, \dots, x_m]$-module and $\mathcal{F}_x$ is the increasing filtration induced by this grading structure. We call the grading of the graded $\mathbb{C}[x_1, \dots, x_m]$-module $C_P(\Gamma)$ the x-grading of $C_P(\Gamma)$. Denote by d_0 and d_1 the two differential maps of the matrix factorization $C_P(\Gamma)$. We decompose d_0 and d_1 into sums of homogeneous components with respect to the x-grading. That is,

$$d_0 = \sum_{l=0}^{\lfloor \frac{N}{k} \rfloor} a^l d_0^{(l)}, \tag{3.5}$$

$$d_1 = \sum_{l=0}^{\lfloor \frac{N}{k} \rfloor} a^l d_1^{(l)}, \tag{3.6}$$

where $d_1^{(l)}$ and $d_0^{(l)}$ are R-module homomorphisms and satisfy:

- they are homogeneous with respect to both the polynomial grading and the x-grading,
- $\deg d_0^{(l)} = \deg d_1^{(l)} = \deg_x d_0^{(l)} = \deg_x d_1^{(l)} = N + 1 - 2lk$.

Recall that $C_P(\Gamma)$ is a matrix factorization of 0. So $d_0 \circ d_1 = 0$ and $d_1 \circ d_0 = 0$. Comparing the homogeneous parts with respect to the x-grading, one gets that, for any $l \geq 0$,

$$\sum_{i=0}^{l} d_0^{(i)} \circ d_1^{(l-i)} = 0, \tag{3.7}$$

$$\sum_{i=0}^{l} d_1^{(i)} \circ d_0^{(l-i)} = 0, \tag{3.8}$$

where we use the convention that $d_0^{(i)} = 0$ and $d_1^{(i)} = 0$ if $i > \lfloor \frac{N}{k} \rfloor$.

By the definition of $C_N(\Gamma)$, there is an isomorphism of $\mathbb{Z}_2$-periodic chain complexes of $\mathbb{C}[a]$-modules

$$C_N(\Gamma) \otimes_{\mathbb{C}} \mathbb{C}[a] \cong C_P^0(\Gamma) \xrightarrow{d_0^{(0)}} C_P^1(\Gamma) \xrightarrow{d_1^{(0)}} C_P^0(\Gamma),$$

that preserves the $\mathbb{Z}_2$-grading, the polynomial grading and the x-grading. So there is an isomorphism of $\mathbb{C}[a]$-modules

$$H_N(\Gamma) \otimes_{\mathbb{C}} \mathbb{C}[a] \cong H(C_P(\Gamma), d^{(0)}) \tag{3.9}$$

preserving the $\mathbb{Z}_2$-grading, the polynomial grading and the x-grading.

Now we are ready to prove that $H_P^{\varepsilon+1}(\Gamma) = 0$. From [4], we know that $H_N^{\varepsilon+1}(\Gamma) = 0$. So $H^{\varepsilon+1}(C_P(\Gamma), d^{(0)}) = 0$. That is, $\mathrm{Im}(d_\varepsilon^{(0)}) = \ker(d_{\varepsilon+1}^{(0)})$. Let α be any element in $\ker d_{\varepsilon+1}$ that is homogeneous with respect to the polynomial grading with $\deg \alpha = g$. Recall that $\deg a = 2k$. So, when we decompose α according to the x-grading, we get $\alpha = \sum_{i=-\infty}^{\infty} \alpha_l$, where α_l is homogeneous with respect to both the polynomial grading and the x-grading with $\deg \alpha_l = g$, $\deg_x \alpha_l = g - 2kl$. Of course, $\alpha_l = 0$ for $l < 0$ and $l \gg 1$ since the x-grading is bounded below.

Next, we construct inductively a sequence $\{\beta_l\}_{-\infty}^{\infty} \subset C_P^\varepsilon(\Gamma)$ such that

1. $\beta_l = 0$ for $l < 0$;
2. each β_l is homogeneous with respect to both the polynomial grading and the x-grading;
3. $\deg \beta_l = g - N - 1$ and $\deg_x \beta_l = g - 2kl - N - 1$;
4. $\alpha_l = \sum_{i=0}^{\lfloor \frac{N}{k} \rfloor} a^i d_\varepsilon^{(i)} \beta_{l-i}$ for all $l \in \mathbb{Z}$.

Again, since the x-grading is bounded below, $\beta_l = 0$ for $l \gg 1$. Note that $\{\beta_l\}_{-\infty}^{-1}$ is the zero sequence and satisfies conditions (1-4) for l up to -1. Now assume that, for some $l \geq 0$, there is a sequence $\{\beta_l\}_{-\infty}^{l-1}$ satisfies conditions (1-4) up to $l - 1$. Let us construct β_l. In the equation $d_{\varepsilon+1}\alpha = 0$, compare the homogeneous parts with respect to the x-grading of x-degree $N+1+g-2lk$. This gives us

$$\begin{aligned}
0 &= \sum_{j=0}^{\lfloor \frac{N}{k} \rfloor} a^j d_{\varepsilon+1}^{(j)} \alpha_{l-j} = d_{\varepsilon+1}^{(0)} \alpha_l + \sum_{j=1}^{\lfloor \frac{N}{k} \rfloor} a^j d_{\varepsilon+1}^{(j)} \alpha_{l-j} \\
&= d_{\varepsilon+1}^{(0)} \alpha_l + \sum_{j=1}^{\lfloor \frac{N}{k} \rfloor} a^j d_{\varepsilon+1}^{(j)} \sum_{i=0}^{\lfloor \frac{N}{k} \rfloor} a^i d_{\varepsilon}^{(i)} \beta_{l-j-i} \\
&= d_{\varepsilon+1}^{(0)} \alpha_l + \sum_{p=1}^{2\lfloor \frac{N}{k} \rfloor} a^p \Big(\sum_{i+j=p,\ 0 \leq i \leq \lfloor \frac{N}{k} \rfloor,\ 1 \leq j \leq \lfloor \frac{N}{k} \rfloor} d_{\varepsilon+1}^{(j)} d_{\varepsilon}^{(i)} \Big) \beta_{l-p} \\
\text{(by (3.7) and (3.8))} \quad &= d_{\varepsilon+1}^{(0)} \alpha_l - \sum_{p=1}^{\lfloor \frac{N}{k} \rfloor} a^p d_{\varepsilon+1}^{(0)} d_{\varepsilon}^{(p)} \beta_{l-p} = d_{\varepsilon+1}^{(0)} \Big(\alpha_l - \sum_{p=1}^{\lfloor \frac{N}{k} \rfloor} a^p d_{\varepsilon}^{(p)} \beta_{l-p} \Big).
\end{aligned}$$

So $\alpha_l - \sum_{p=1}^{\lfloor \frac{N}{k} \rfloor} a^p d_{\varepsilon}^{(p)} \beta_{l-p} \in \ker(d_{\varepsilon+1}^{(0)}) = \mathrm{Im}(d_{\varepsilon}^{(0)})$. Therefore, there is a β_l satisfying conditions (1-2) such that $d_{\varepsilon}^{(0)} \beta_l = \alpha_l - \sum_{p=1}^{\lfloor \frac{N}{k} \rfloor} a^p d_{\varepsilon}^{(p)} \beta_{l-p}$. Thus, $\{\beta_l\}_{-\infty}^{l}$ satisfies conditions (1-4) above. This completes the inductive construction. Note that $\sum_{l=-\infty}^{\infty} \beta_l$ is a finite sum and therefore a well defined element of $C_P^{\varepsilon}(\Gamma)$. We have

$$\begin{aligned}
\alpha &= \sum_{l=-\infty}^{\infty} \alpha_l = \sum_{l=-\infty}^{\infty} \sum_{i=0}^{\lfloor \frac{N}{k} \rfloor} a^i d_{\varepsilon}^{(i)} \beta_{l-i} \\
&= \sum_{i=0}^{\lfloor \frac{N}{k} \rfloor} a^i d_{\varepsilon}^{(i)} \Big(\sum_{l=-\infty}^{\infty} \beta_{l-i} \Big) = \sum_{i=0}^{\lfloor \frac{N}{k} \rfloor} a^i d_{\varepsilon}^{(i)} \Big(\sum_{l=-\infty}^{\infty} \beta_l \Big) \\
&= d_{\varepsilon} \Big(\sum_{l=-\infty}^{\infty} \beta_l \Big).
\end{aligned}$$

So $\alpha \in \mathrm{Im} d_{\varepsilon}$. This shows $\mathrm{Im}(d_{\varepsilon}) = \ker(d_{\varepsilon+1})$ and therefore $H_P^{\varepsilon+1}(\Gamma) = 0$.

It remains to prove (3.4). According to (3.9), we only need to show that

$$\mathcal{F}_x^n H_P^{\varepsilon}(\Gamma) / \mathcal{F}_x^{n-1} H_P^{\varepsilon}(\Gamma) \cong H^{\varepsilon,n}(C_P(\Gamma), d^{(0)}), \tag{3.10}$$

where $H^{\varepsilon,n}(C_P(\Gamma), d^{(0)})$ is the direct sum component of the free $\mathbb{C}[a]$-module $H(C_P(\Gamma), d^{(0)})$ consisting of homogeneous elements of $\mathbb{Z}_2$-grading ε and x-grading n.

Denote by $(\ker d_{\varepsilon}^{(0)})_n$ the $\mathbb{C}[a]$-submodule of $\ker d_{\varepsilon}^{(0)}$ consisting of elements homogeneous with respective to the x-grading of x-degree n. Next, for every $\alpha \in (\ker d_{\varepsilon}^{(0)})_n$, we construct by induction a sequence $\{\alpha_l\}_0^{\infty} \subset C_P^{\varepsilon}(\Gamma)$, such that $\alpha_0 = \alpha$, α_l is a homogeneous element with respective to the x-grading of x-degree $n - 2lk$, and

$$\sum_{j=0}^{l} a^j d_{\varepsilon}^{(j)} \alpha_{l-j} = 0, \ \forall \ l \in \mathbb{Z}, \tag{3.11}$$

where we use the convention that $d_{\varepsilon}^{(j)} = 0$ for $j > \lfloor \frac{N}{k} \rfloor$. Again, since the x-grading is bounded below, $\alpha_l = 0$ for $l \gg 1$. Clearly, $\{\alpha_l\}_0^0$ with $\alpha_0 = \alpha$ satisfies the initial condition and equation (3.11) up to $l = 0$. Assume that, for some $l \geq 1$, $\{\alpha_l\}_0^{l-1}$ is constructed and satisfies the initial condition and equation (3.11) up to $l - 1$.

Let us find an α_l. Note that

$$\begin{aligned}
d^{(0)}_{\varepsilon+1}(\sum_{j=1}^{l} a^j d^{(j)}_{\varepsilon} \alpha_{l-j}) &= \sum_{j=1}^{l} a^j d^{(0)}_{\varepsilon+1} d^{(j)}_{\varepsilon} \alpha_{l-j} \\
\text{(by (3.7) and (3.8))} &= -\sum_{j=1}^{l} a^j \sum_{i=0}^{j-1} d^{(j-i)}_{\varepsilon+1} d^{(i)}_{\varepsilon} \alpha_{l-j} \\
\text{(setting } p = l-j+i\text{)} &= -\sum_{p=0}^{l-1} a^{l-p} \sum_{i=0}^{p} a^i d^{(l-p)}_{\varepsilon+1} d^{(i)}_{\varepsilon} \alpha_{p-i} \\
&= -\sum_{p=0}^{l-1} a^{l-p} d^{(l-p)}_{\varepsilon+1} (\sum_{i=0}^{p} a^i d^{(i)}_{\varepsilon} \alpha_{p-i}) \\
\text{(by induction hypothesis)} &= 0
\end{aligned}$$

But $H^{\varepsilon+1}(C_P(\Gamma), d^{(0)}) = 0$, that is, $\mathrm{Im}(d^{(0)}_{\varepsilon}) = \ker(d^{(0)}_{\varepsilon+1})$. So there is an $\alpha_l \in C^{\varepsilon}_P(\Gamma)$ homogeneous with respective to the x-grading of x-degree $n - 2lk$ satisfying $d^{(0)}_{\varepsilon}\alpha_l = -\sum_{j=1}^{l} a^j d^{(j)}_{\varepsilon} \alpha_{l-j}$. Thus, the sequence $\{\alpha_l\}_0^l$ satisfies the initial condition and equation (3.11) up to l. This completes the induction and we have the sequence $\{\alpha_l\}_0^\infty$. As explained above, $\sum_{l=0}^\infty \alpha_l$ is in fact a finite sum and therefore a well defined element of $C^{\varepsilon}_P(\Gamma)$. Note that the homogeneous part of $d_\varepsilon(\sum_{l=0}^\infty \alpha_l)$ with respect to the x-grading of x-degree $N + 1 + n - 2lk$ is $\sum_{j=0}^{l} a^j d^{(j)}_{\varepsilon} \alpha_{l-j} = 0$ by equation (3.11). This implies that $d_\varepsilon(\sum_{l=0}^\infty \alpha_l) = 0$, that is, $\sum_{l=0}^\infty \alpha_l$ is a cycle in $(C_P(\Gamma), d)$.

Define $\tilde{\phi}_n : (\ker d^{(0)}_{\varepsilon})_n \to \mathcal{F}^n_x H^{\varepsilon}_P(\Gamma)/\mathcal{F}^{n-1}_x H^{\varepsilon}_P(\Gamma)$ by $\alpha \mapsto [\sum_{l=0}^\infty \alpha_l]$. Since the top homogeneous component of $\sum_{l=0}^\infty \alpha_l$ with respect to the x-grading is $\alpha_0 = \alpha$, one can see that $\tilde{\phi}_n(\alpha)$ does not depend on the choice of $\{\alpha_l\}_0^\infty$ and is well defined. It is also easy to verify that $\tilde{\phi}_n$ is a $\mathbb{C}[a]$-module homomorphism preserving the polynomial grading. Moreover, $\tilde{\phi}_n$ is surjective. To see this, note that any element of $\mathcal{F}^n_x H^{\varepsilon}_P(\Gamma)/\mathcal{F}^{n-1}_x H^{\varepsilon}_P(\Gamma)$ can be expressed as $[\sum_{l=0}^\infty \alpha_l]$, where α_l is a homogeneous element with respective to the x-grading of x-degree $n - 2lk$ and $d_\varepsilon \sum_{l=0}^\infty \alpha_l = 0$. Comparing the top homogeneous parts with respect to the x-grading on both sides of this equation, one gets $d^{(0)}_{\varepsilon}\alpha_0 = 0$, which means $\alpha_0 \in (\ker d^{(0)}_{\varepsilon})_n$. By the definition of $\tilde{\phi}_n$, one easily sees that $\tilde{\phi}_n(\alpha_0) = [\sum_{l=0}^\infty \alpha_l]$.

Denote by $(\mathrm{Im} d^{(0)}_{\varepsilon+1})_n$ the homogeneous component of $\mathrm{Im} d^{(0)}_{\varepsilon+1}$ with respect to the x-grading of x-degree n. We prove isomorphism (3.10) by showing that $\ker \tilde{\phi}_n = (\mathrm{Im} d^{(0)}_{\varepsilon+1})_n$. Assume $\alpha \in \ker \tilde{\phi}_n$ and $\{\alpha_l\}_0^\infty$ is a sequence given by the above inductive construction. Then

$$\sum_{l=0}^{\infty} \alpha_l = d_{\varepsilon+1}\beta + \gamma, \tag{3.12}$$

where γ is a cycle in $\mathcal{F}^{n-1}_x C^{\varepsilon}_P(\Gamma)$, and $\beta \in C^{\varepsilon+1}_P(\Gamma)$ satisfying $d_{\varepsilon+1}\beta \in \mathcal{F}^n_x C^{\varepsilon}_P(\Gamma)$. We claim that we can choose β so that $\deg_x \beta \leq n - N - 1$. Assume that $\deg_x \beta = g > n - N - 1$ and denote by β_0 the top homogeneous part of β with respect to the x-grading. Comparing the top homogeneous parts with respect to the x-grading on both sides of equation (3.12), we have $d^{(0)}_{\varepsilon+1}\beta_0 = 0$. So there exists a homogeneous element $\theta \in C^{\varepsilon}_P(\Gamma)$ of degree $g - N - 1$ such that $d^{(0)}_{\varepsilon}\theta = \beta_0$. Let $\beta' = \beta - d_\varepsilon\theta$. Then β' also satisfies the above equation, and $\deg_x \beta' < \deg_x \beta$. Repeat this process. Within finite steps, we get a $\hat{\beta} \in C^{\varepsilon+1}_P(\Gamma)$ with $\deg_x \hat{\beta} \leq n - N - 1$ and

$$\sum_{l=0}^{\infty} \alpha_l = d_{\varepsilon+1}\hat{\beta} + \gamma. \tag{3.13}$$

Let $\hat{\beta}_0$ be the homogeneous part of $\hat{\beta}$ with respect to the x-grading of x-degree $n - N - 1$. Comparing the top homogeneous parts with respect to the x-grading on both sides of equation (3.13), one can see that $\alpha = \alpha_0 = d^{(0)}_{\varepsilon+1}\hat{\beta}_0$. This shows $\alpha \in (\mathrm{Im} d^{(0)}_{\varepsilon+1})_n$. So $\ker \tilde{\phi}_n \subset (\mathrm{Im} d^{(0)}_{\varepsilon+1})_n$. On the other hand, if $\alpha \in (\mathrm{Im} d^{(0)}_{\varepsilon+1})_n$, then $\alpha = d^{(0)}_{\varepsilon+1}\beta$ for some $\beta \in C_P^{\varepsilon+1}(\Gamma)$ homogeneous with respect to the x-grading of x-degree $n-N-1$. So

$$\sum_{l=0}^{\infty} \alpha_l = d^{(0)}_{\varepsilon+1}\beta + \sum_{l=1}^{\infty} \alpha_l = d_{\varepsilon+1}\beta + (\sum_{l=1}^{\infty} \alpha_l - \sum_{j=1}^{\lfloor \frac{N}{k} \rfloor} a^j d^{(j)}_{\varepsilon+1}\beta) \in \ker \tilde{\phi}_n.$$

Thus, $(\mathrm{Im} d^{(0)}_{\varepsilon+1})_n \subset \ker \tilde{\phi}_n$. This shows $(\mathrm{Im} d^{(0)}_{\varepsilon+1})_n = \ker \tilde{\phi}_n$ and, therefore, $\tilde{\phi}_n$ induces a $\mathbb{C}[a]$-module isomorphism $\phi_n : H^{\varepsilon,n}(C_P(\Gamma), d^{(0)}) \to \mathcal{F}^n_x H^{\varepsilon}_P(\Gamma)/\mathcal{F}^{n-1}_x H^{\varepsilon}_P(\Gamma)$ preserving the polynomial grading. ☐

Corollary 3.3. *Let* Γ *be a closed MOY graph. Then*

(1) $\hat{H}_P(\Gamma)$ *is a finite dimensional* $\mathbb{C}$*-space and its* x*-filtration is bounded and exhaustive.*
(2) $H_P(\Gamma)$ *is a finitely generated graded-free* $\mathbb{C}[a]$*-module and its* x*-filtration is bounded and exhaustive, where "graded-free" means* $H_P(\Gamma)$ *is graded, free and admits a homogeneous basis.*

Proof. From [4], we know that $H_N(\Gamma)$ is finite dimensional and its polynomial grading is bound above and below. In addition, by their definitions, we know that the x-filtrations of $\hat{H}_P(\Gamma)$ and $H_P(\Gamma)$ are bounded below and exhaustive. Using Lemmas 3.1 and 3.2, one can inductively prove that, for every n,

- $\mathcal{F}^n_x \hat{H}_P(\Gamma)$ is a finite dimensional $\mathbb{C}$-space,
- $\mathcal{F}^n_x H_P(\Gamma)$ is a finitely generated free $\mathbb{C}[a]$-module.

Lemmas 3.1 and 3.2 also imply that the x-filtrations of $\hat{H}_P(\Gamma)$ and $H_P(\Gamma)$ are bounded above. Thus,

- $\hat{H}_P(\Gamma)$ is itself a finite dimensional $\mathbb{C}$-space,
- $H_P(\Gamma)$ is itself a finitely generated free $\mathbb{C}[a]$-module.

Finally, since the polynomial grading of $H_P(\Gamma)$ is bounded below, we know that $H_P(\Gamma)$ is a graded-free $\mathbb{C}[a]$-module by, for example, [12, Lemma 3.3]. ☐

3.2. $E_r(L)$ **and** $\hat{E}_r(L)$**.** Using Lemmas 3.1, 3.2 and Corollary 3.3, it is straightforward to prove Theorem 1.5. We summarize the key observation in the proof as the following lemma.

Lemma 3.4. *Suppose* Γ_1 *is a closed MOY graph and* Γ_0 *is obtained from* Γ_1 *by replacing a wide edge by a pair of parallel regular edges.*[7] *Denote by* $C_P(\Gamma_0) \underset{\chi_1}{\overset{\chi_0}{\rightleftarrows}} C_P(\Gamma_1)$ *the homomorphisms induced by this local change and by* $\chi_0^{(0)}$, $\chi_1^{(0)}$ *the top homogeneous parts of* χ_0, χ_1 *with respect to the* x*-grading. In addition, we denote by* $d^{(0)}_{mf}$ *the top homogeneous parts of the differential maps of* $C_P(\Gamma_0)$ *and* $C_P(\Gamma_1)$ *with respect to the* x*-grading. Then*

- $(C_P(\Gamma_0), d^{(0)}_{mf}) \underset{\chi_1^{(0)}}{\overset{\chi_0^{(0)}}{\rightleftarrows}} (C_P(\Gamma_0), d^{(0)}_{mf})$ *are homomorphisms of matrix factorizations of* 0.
- *The following squares commute, where* ε *is the rotation number of* Γ_0 *and* Γ_1*, and* ϕ_{n,Γ_0}*,* ϕ_{n,Γ_1} *are the isomorphisms constructed in the proof of Lemma 3.2.*

$$\begin{tikzcd}
H^{\varepsilon,n}(C_P(\Gamma_0), d^{(0)}_{mf}) \arrow[r, "\phi_{n,\Gamma_0}"] \arrow[d, "\chi_0^{(0)}"] & \mathcal{F}^n_x H^{\varepsilon}_P(\Gamma_0)/\mathcal{F}^{n-1}_x H^{\varepsilon}_P(\Gamma_0) \arrow[d, "\chi_0"] \\
H^{\varepsilon,n}(C_P(\Gamma_1), d^{(0)}_{mf}) \arrow[r, "\phi_{n,\Gamma_1}"] & \mathcal{F}^n_x H^{\varepsilon}_P(\Gamma_1)/\mathcal{F}^{n-1}_x H^{\varepsilon}_P(\Gamma_1)
\end{tikzcd}
\qquad
\begin{tikzcd}
H^{\varepsilon,n}(C_P(\Gamma_1), d^{(0)}_{mf}) \arrow[r, "\phi_{n,\Gamma_1}"] \arrow[d, "\chi_1^{(0)}"] & \mathcal{F}^n_x H^{\varepsilon}_P(\Gamma_1)/\mathcal{F}^{n-1}_x H^{\varepsilon}_P(\Gamma_1) \arrow[d, "\chi_1"] \\
H^{\varepsilon,n}(C_P(\Gamma_0), d^{(0)}_{mf}) \arrow[r, "\phi_{n,\Gamma_0}"] & \mathcal{F}^n_x H^{\varepsilon}_P(\Gamma_0)/\mathcal{F}^{n-1}_x H^{\varepsilon}_P(\Gamma_0)
\end{tikzcd}$$

Proof. This lemma follows easily from the constructions of χ_0, χ_1 and ϕ_n. We leave the details to the reader. ☐

It is now very easy to prove Theorem 1.5.

---

[7] That is, replacing a piece of Γ_1 of the form $\Gamma_{i,j;p,q}$ in Figure 3 by $\Gamma_{i;p} \sqcup \Gamma_{j;q}$ in the same figure.

Proof of Theorem 1.5. By Corollary 3.3, the x-filtrations of $H(\hat{C}_P(D), d_{mf})$ and $H(C_P(D), d_{mf})$ are bounded and exhaustive. So $\hat{E}_r(L)$ and $E_r(L)$ converge to $\hat{H}_P(L)$ and $H_P(L)$, respectively. It remains to show that $\hat{E}_1(L) \cong H_N(L)$ and $E_1(L) \cong H_N(L) \otimes_{\mathbb{C}} \mathbb{C}[a]$. The proofs of these isomorphisms are very similar, and the proof of $\hat{E}_1(L) \cong H_N(L)$ can be found in [2, 11]. So we only prove $E_1(L) \cong H_N(L) \otimes_{\mathbb{C}} \mathbb{C}[a]$ here. By Lemma 3.4, we know that $E_0(L)$ is isomorphic to the chain complex $(H(C_P(D), d_{mf}^{(0)}), d_\chi^{(0)})$, where $d_{mf}^{(0)}$ and $d_\chi^{(0)}$ are the top homogeneous parts of d_{mf} and d_χ with respect to the x-grading of $C_P(D)$. So $E_1(L) \cong H(H(C_P(D), d_{mf}^{(0)}), d_\chi^{(0)})$. On the other hand, by the definition of $H_N(L)$, it is easy to see that $H(H(C_P(D), d_{mf}^{(0)}), d_\chi^{(0)}) \cong H_N(L) \otimes_{\mathbb{C}} \mathbb{C}[a]$. So $E_1(L) \cong H_N(L) \otimes_{\mathbb{C}} \mathbb{C}[a]$. □

4. Decomposition Theorems

In this section, we prove Theorems 1.7, 1.11 and explicitly compute the spectral sequences of the filtered chain complexes $F_{i,s}$, $T_{i,m,s}$, $\hat{F}_{i,s}$ and $\hat{T}_{i,m,s}$ defined in (1.9)–(1.12).

4.1. A closer look at $\mathcal{F}_x$.

To prove Theorems 1.7 and 1.11, we need to better understand the relation between the polynomial grading and the x-filtration. The goal of this subsection is to show that, for a closed MOY graph Γ, any direct sum decomposition of $H_P(\Gamma)$ in the category of graded $\mathbb{C}[a]$-modules is also a direct sum decomposition in the category of filtered $\mathbb{C}$-spaces. Theorems 1.7 and 1.11 both follow from this.

In the rest of this subsection, Γ is a closed MOY graph with a marking, $x_1, \dots, x_m$ are the variables assigned to the marked points of Γ and $R = \mathbb{C}[x_1, \dots, x_m, a]$, where a is a homogeneous variable of degree $2k$. $P(x, a)$ is the homogeneous polynomial given in (1.1). Unless otherwise specified, when we say an element is homogeneous, we mean it is homogeneous with respect to the polynomial grading.

We start with simple observations.

Lemma 4.1. *Let M be a finitely generated free $\mathbb{C}[a]$-module and $\{v_i\}$ a basis for M. For any $\lambda \in \mathbb{C}$, denote by $\pi_\lambda : M \to M/(a-\lambda)M$ the standard quotient map. Then $\{\pi_\lambda(v_i)\}$ is a basis for the $\mathbb{C}$-space $M/(a-\lambda)M$.*

Proof. Since $\{v_i\}$ spans M, we know that $\{\pi_\lambda(v_i)\}$ spans $M/(a-\lambda)M$. It remains to show that $\{\pi_\lambda(v_i)\}$ is linearly independent. Suppose $\{c_i\} \subset \mathbb{C}$ satisfies $\sum_i c_i \pi_\lambda(v_i) = 0$. Then $\sum_i c_i v_i \in (a-\lambda)M$. Therefore, there are $f_i \in R$ such that $\sum_i c_i v_i = (a-\lambda)\sum_i f_i v_i$. That is, $\sum_i (c_i - (a-\lambda)f_i)v_i = 0$. Since $\{v_i\}$ a basis for M, this means $c_i - (a-\lambda)f_i$ and, therefore, $c_i = f_i = 0$ for every i. □

Lemma 4.2. *For $f \in R$, denote by f_i the homogeneous component of f with $\deg f_i = i$. Then $\deg_x f_i \leq \deg_x f$ for every i.*

Proof. Obvious. □

Lemma 4.3. *(1) If $u \in \mathcal{F}_x^n C_P(\Gamma)$, then all homogeneous components of u are also in $\mathcal{F}_x^n C_P(\Gamma)$.*
(2) If $[u] \in \mathcal{F}_x^n H_P(\Gamma)$, then all homogeneous components of $[u]$ are also in $\mathcal{F}_x^n H_P(\Gamma)$.

Proof. $C_P(\Gamma)$ is a Koszul matrix factorization. Denote by $\{1_{\vec{\varepsilon}}\}$ the standard basis for $C_P(\Gamma)$ given in Definition 2.4. Recall that $\{1_{\vec{\varepsilon}}\}$ is a homogeneous basis with respect to the polynomial grading.

To prove Part (1) of the lemma, assume $u \in \mathcal{F}_x^n C_P(\Gamma)$ and denote by u_i the homogeneous component of u with $\deg u_i = i$. Every u_i can be uniquely expressed as $u_i = \sum_{\vec{\varepsilon}} g_{i,\vec{\varepsilon}} 1_{\vec{\varepsilon}}$, where $g_{i,\vec{\varepsilon}}$ is a homogeneous element of R with $\deg g_{i,\vec{\varepsilon}} = i - \deg 1_{\vec{\varepsilon}}$. Then $u = \sum_i u_i = \sum_{\vec{\varepsilon}} (\sum_i g_{i,\vec{\varepsilon}}) 1_{\vec{\varepsilon}}$. Since $u \in \mathcal{F}_x^n C_P(\Gamma)$, we have $\deg_x \sum_i g_{i,\vec{\varepsilon}} \leq n - \deg_x 1_{\vec{\varepsilon}}$ for every $\vec{\varepsilon}$. Note that $g_{i,\vec{\varepsilon}}$ is the homogeneous component of $\sum_i g_{i,\vec{\varepsilon}}$ of polynomial degree $i - \deg 1_{\vec{\varepsilon}}$. Thus, by Lemma 4.2, we have $\deg_x g_{i,\vec{\varepsilon}} \leq n - \deg_x 1_{\vec{\varepsilon}}$. So $\deg_x u_i = \deg_x \sum_{\vec{\varepsilon}} g_{i,\vec{\varepsilon}} 1_{\vec{\varepsilon}} \leq n$. This proves Part (1).

To prove Part (2), note that $[u]$ is represented by a cycle $u \in \mathcal{F}_x^n C_P(\Gamma)$. Denote by u_i the homogeneous component of u with $\deg u_i = i$. Since the differential of $C_P(\Gamma)$ is homogeneous, each u_i is a cycle, and $[u_i]$ is the homogeneous component of $[u]$ with $\deg [u_i] = i$. Then Part (2) of the lemma follows from Part (1). □

By Lemma 3.3, $H_P(\Gamma)$ is a finitely generated graded-free $\mathbb{C}[a]$-module. The next lemma determines the x-filtration degrees of elements of homogeneous bases for $H_P(\Gamma)$.

Lemma 4.4. *Let $\{[u_j]\}$ be any homogeneous basis for the free $\mathbb{C}[a]$-module $H_P(\Gamma)$. Then $\deg_x [u_j] = \deg [u_j]$ for every j.*

Proof. By Lemma 2.5, we know that $\deg_x[u_j] \leq \deg[u_j]$ for every j. Assume $n = \deg_x[u_j] < \deg[u_j] = l$ for a certain j. Then $[u_j]$ is represented by a cycle $u_j \in \mathcal{F}_x^n C_P(\Gamma)$. Denote by $u_{j,i}$ the homogeneous component of u_j with $\deg u_{j,i} = i$. By Lemma 4.3, we know that $u_{j,i} \in \mathcal{F}_x^n C_P(\Gamma)$ for every i. Also, since the differential of $C_P(\Gamma)$ is homogeneous, each $u_{j,i}$ is itself a cycle. By comparing the homogeneous components in $[u_j] = \sum_i [u_{j,i}]$, we get $[u_j] = [u_{j,l}]$ and $[u_{j,i}] = 0$ if $i \neq l$. Note that $\deg_x u_{j,l} \leq n < l = \deg u_{j,l}$. So, by Lemma 2.5, $u_{j,l} = av$ for some $v \in C_P(\Gamma)$. It is easy to see that v is a homogeneous cycle in $C_P(\Gamma)$ and that $\{[v]\} \cup \{u_i \mid i \neq j\}$ spans $H_P(\Gamma)$ and is $\mathbb{C}[a]$-linearly independent. In other words, $\{[v]\} \cup \{[u_i] \mid i \neq j\}$ is also a basis for $H_P(\Gamma)$. But this is impossible because, if this is true, then the determinant of the change-of-coordinates matrix from the basis $\{[u_j]\}$ to the basis $\{[v]\} \cup \{[u_i] \mid i \neq j\}$ is a, which is not invertible in $\mathbb{C}[a]$. □

Definition 4.5. Denote by $\pi_0 : C_P(\Gamma) \to C_N(\Gamma)(= C_P(\Gamma)/aC_P(\Gamma))$ the standard quotient map. To keep notations simple, we denote again by $\pi_0 : H_P(\Gamma) \to H_N(\Gamma)$ the homomorphism induced by the quotient map π_0.

Recall that $C_N(\Gamma)$ inherits the polynomial grading of $C_P(\Gamma)$ via π_0, which makes π_0 a homogeneous map of degree 0. Moreover, $C_N(\Gamma)$ also inherits the x-filtration of $C_P(\Gamma)$ via π_0. It is easy to see the x-filtration of $C_N(\Gamma)$ is the filtration induced by its polynomial grading.

Lemma 4.6. *$\pi_0 : H_P(\Gamma) \to H_N(\Gamma)$ is a surjective homogeneous homomorphism with $\deg \pi_0 = 0$ and $\ker \pi_0 = aH_P(\Gamma)$. Moreover, any homogeneous basis for the free $\mathbb{C}[a]$-module $H_P(\Gamma)$ is mapped by π_0 to a homogeneous basis for the $\mathbb{C}$-space $H_N(\Gamma)$.*

Proof. By its definition, we know that $\pi_0 : H_P(\Gamma) \to H_N(\Gamma)$ is a homogeneous homomorphism with $\deg \pi_0 = 0$. The short exact sequence $0 \to C_P(\Gamma) \xrightarrow{a} C_P(\Gamma) \xrightarrow{\pi_0} C_N(\Gamma) \to 0$ induces a long exact sequence

$$\cdots \to H_N^{\varepsilon+1}(\Gamma) \to H_P^{\varepsilon}(\Gamma) \xrightarrow{a} H_P^{\varepsilon}(\Gamma) \xrightarrow{\pi_0} H_N^{\varepsilon}(\Gamma) \to H_P^{\varepsilon+1}(\Gamma) \xrightarrow{a} H_P^{\varepsilon+1}(\Gamma) \xrightarrow{\pi_0} H_N^{\varepsilon+1}(\Gamma) \to \cdots$$

with $\mathbb{Z}_2$ homological grading, where ε is the rotation number of Γ. From [4], we know that $H_N^{\varepsilon+1}(\Gamma) \cong 0$. By Lemma 3.2, we know that $H_P^{\varepsilon+1}(\Gamma) \cong 0$. So the above long exact sequence becomes a short exact sequence

$$0 \to H_P(\Gamma) \xrightarrow{a} H_P(\Gamma) \xrightarrow{\pi_0} H_N(\Gamma) \to 0.$$

It follows from this that $\pi_0 : H_P(\Gamma) \to H_N(\Gamma)$ is surjective and $\ker \pi_0 = aH_P(\Gamma)$. The statement about bases follows then from Lemma 4.1. □

Lemma 4.7. *For any $[u] \in H_P(\Gamma)$, $[u] \in \mathcal{F}_x^n H_P(\Gamma)$ if and only if $a[u] \in \mathcal{F}_x^n H_P(\Gamma)$.*

Proof. Since the map $C_P(\Gamma) \xrightarrow{a} C_P(\Gamma)$ preserves the x-filtration, so does the map $H_P(\Gamma) \xrightarrow{a} H_P(\Gamma)$. Therefor, $a[u] \in \mathcal{F}_x^n H_P(\Gamma)$ if $[u] \in \mathcal{F}_x^n H_P(\Gamma)$.

Now assume $[u] \notin \mathcal{F}_x^n H_P(\Gamma)$. Then $[u] \neq 0$. Since the x-filtration is exhaustive, there is an $l > n$ such that $[u] \in \mathcal{F}_x^l H_P(\Gamma)$ and $[u] \notin \mathcal{F}_x^{l-1} H_P(\Gamma)$. Recall that, by Lemma 3.2, we have $H_P^{\varepsilon+1}(\Gamma) = 0$. Furthermore, in the proof of Lemma 3.2, we constructed an isomorphism $\phi_l : H_N^{\varepsilon,l}(\Gamma) \otimes_{\mathbb{C}} \mathbb{C}[a] \to \mathcal{F}_x^l H_P^{\varepsilon}(\Gamma)/\mathcal{F}_x^{l-1} H_P^{\varepsilon}(\Gamma)$ of $\mathbb{C}[a]$-modules. So $[u] \in H_P^{\varepsilon}(\Gamma)$ and $\phi_l^{-1}([u]) \neq 0$. But $H_N^{\varepsilon,l}(\Gamma) \otimes_{\mathbb{C}} \mathbb{C}[a]$ is a free $\mathbb{C}[a]$-module. So $\phi_l^{-1}(a[u]) = a\phi_l^{-1}([u]) \neq 0$. Thus, $a[u] \notin \mathcal{F}_x^{l-1} H_P(\Gamma)$ and, therefore, $a[u] \notin \mathcal{F}_x^n H_P(\Gamma)$. □

Lemma 4.8. *Let $\{[u_j]\}$ be a homogeneous basis for $H_P(\Gamma)$. For any $\{f_j\} \subset \mathbb{C}[a]$ and $l \in \mathbb{Z}$, $\sum_j f_j[u_j] \in \mathcal{F}_x^l H_P(\Gamma)$ if and only if $f_j = 0$ whenever $\deg[u_j] > l$.*

Proof. By Lemma 4.4, we have $\deg_x[u_j] = \deg[u_j]$ for every j. If $f_j = 0$ whenever $\deg[u_j] > l$, then, by Lemma 4.7, we know that $\sum_j f_j[u_j] \in \mathcal{F}_x^l H_P(\Gamma)$.

Now assume $\sum_j f_j[u_j] \in \mathcal{F}_x^l H_P(\Gamma)$. We prove by contradiction that $f_j = 0$ whenever $\deg[u_j] > l$. Assuming the conclusion is not true, then $n := \{\deg[u_j] \mid f_j \neq 0\} > l$. Without loss of generality, we assume

$$n_j := \deg_x[u_j] = \deg[u_j] \begin{cases} = n & \text{if } 1 \leq j \leq p, \\ < n & \text{if } p+1 \leq j \leq p+q, \\ > n & \text{otherwise.} \end{cases}$$

By the definition of n, we know that $f_j = 0$ unless $1 \leq j \leq p+q$. For $1 \leq j \leq p+q$, write $f_j = \sum_{i \geq 0} c_{j,i} a^i$. Define $[v_i] = \sum_{j=1}^{p} c_{j,i}[u_j] + \sum_{j=p+1}^{p+q} c_{j,n+i-n_j} a^{n-n_j}[u_j]$. Then the homogeneous component of $\sum_j f_j[u_j]$ of polynomial degree $n+i$ is $a^i[v_i]$. By Lemma 4.3, we have $a^i[v_i] \in \mathcal{F}_x^l H_P(\Gamma)$. Therefore, by Lemma 4.7, we have $[v_i] \in \mathcal{F}_x^l H_P(\Gamma)$. Consider $\pi_0([v_i]) = \sum_{j=1}^{p} c_{j,i}\pi_0([u_j]) \in H_N(\Gamma)$. On the one hand, we have that $\pi_0([v_i]) \in \mathcal{F}_x^l H_N(\Gamma)$. On the other hand, we know that $\deg \pi_0([u_1]) = \cdots = \pi_0([u_p]) = n > l$. But the x-filtration on $H_N(\Gamma)$ is induced by the polynomial grading. We must have $\pi_0([v_i]) = \sum_{j=1}^{p} c_{j,i}\pi_0([u_j]) = 0$. Lemma 4.6 tells us that $\{\pi_0([u_1]), \ldots, \pi_0([u_p])\}$ is linearly independent. So $c_{j,i} = 0$ for $i \geq 0$ and $1 \leq j \leq p$. In other words, $f_j = 0$ for $1 \leq j \leq p$. This contradicts the definition of n. ☐

Definition 4.9. Denote by $\pi_1 : C_P(\Gamma) \to \hat{C}_P(\Gamma)(= C_P(\Gamma)/(a-1)C_P(\Gamma))$ the standard quotient map. To keep notations simple, we denote again by $\pi_1 : H_P(\Gamma) \to \hat{H}_P(\Gamma)$ the homomorphism induced by the quotient map π_1.

Note that $\hat{C}_P(\Gamma)$ inherits the x-filtration of $C_P(\Gamma)$ through π_1.

Lemma 4.10. *$\pi_1 : H_P(\Gamma) \to \hat{H}_P(\Gamma)$ is a surjective homomorphism preserving the x-filtration with* $\ker \pi_1 = (a-1)H_P(\Gamma)$*. Moreover, for any homogeneous basis $\{[u_j]\}$ for the free $\mathbb{C}[a]$-module $H_P(\Gamma)$, we have:*

- *$\{\pi_1([u_j])\}$ is a basis for the $\mathbb{C}$-space $\hat{H}_P(\Gamma)$.*
- *$\deg_x \pi_1([u_j]) = \deg[u_j]$ for every j.*
- *For any $\{c_j\} \subset \mathbb{C}$, $\sum_j c_j\pi_1([u_j]) \in \mathcal{F}_x^l \hat{H}_P(\Gamma)$ if and only if $c_j = 0$ whenever $\deg[u_j] > l$.*

Proof. By its definition, we know that $\pi_1 : H_P(\Gamma) \to \hat{H}_P(\Gamma)$ preserves the x-filtration. The short exact sequence $0 \to C_P(\Gamma) \xrightarrow{a-1} C_P(\Gamma) \xrightarrow{\pi_1} \hat{C}_P(\Gamma) \to 0$ induces a long exact sequence

$$\cdots \to \hat{H}_P^{\varepsilon+1}(\Gamma) \to H_P^{\varepsilon}(\Gamma) \xrightarrow{a-1} H_P^{\varepsilon}(\Gamma) \xrightarrow{\pi_1} \hat{H}_P^{\varepsilon}(\Gamma) \to H_P^{\varepsilon+1}(\Gamma) \xrightarrow{a-1} H_P^{\varepsilon+1}(\Gamma) \xrightarrow{\pi_1} \hat{H}_P^{\varepsilon+1}(\Gamma) \to \cdots$$

with $\mathbb{Z}_2$ homological grading, where ε is the rotation number of Γ. By Lemmas 3.1 and 3.2, we know that $\hat{H}_P^{\varepsilon+1}(\Gamma) \cong 0$ and $H_P^{\varepsilon+1}(\Gamma) \cong 0$. So the above long exact sequence becomes a short exact sequence

$$0 \to H_P(\Gamma) \xrightarrow{a-1} H_P(\Gamma) \xrightarrow{\pi_1} \hat{H}_P(\Gamma) \to 0.$$

Thus, $\pi_1 : H_P(\Gamma) \to \hat{H}_P(\Gamma)$ is surjective and $\ker \pi_1 = (a-1)H_P(\Gamma)$.

Now assume $\{[u_j]\}$ is a homogeneous basis for the free $\mathbb{C}[a]$-module $H_P(\Gamma)$ with $\deg[u_j] = n_j$. It follows from the above and Lemma 4.1 that $\{\pi_1([u_j])\}$ is a basis for the $\mathbb{C}$-space $\hat{H}_P(\Gamma)$.

Since $\pi_1 : H_P(\Gamma) \to \hat{H}_P(\Gamma)$ preserves the x-filtration, we get from Lemma 4.4 that $\deg_x \pi_1([u_j]) \leq \deg_x[u_j] = \deg[u_j] = n_j$. Next, we prove that $\deg_x \pi_1([u_j]) = n_j$. Note that $[u_j]$ is represented by a homogeneous cycle u_j in $C_P(\Gamma)$ with $\deg u_j = n_j$. Recall that the x-filtration on $C_P(\Gamma)$ is the increasing filtration associated to an x-grading on $C_P(\Gamma)$. Denote by $u_{j,i}$ the homogeneous component of u_j with respect to this x-grading. Then $u_{j,i} = 0$ if $i > n_j$ and, by Lemma 2.5, $u_{j,i} \in aC_P(\Gamma)$ if $i < n_j$. So $\pi_0(u_{j,n_j}) = \pi_0(u_j)$ is a homogeneous cycle in $C_N(\Gamma)$ of polynomial degree n_j representing the homology class $\pi_0([u_j])$. Lemma 4.6 implies that $\pi_0([u_j]) \neq 0$. Recall that Lemma 3.1 is proved in [11] by a construction very similar to the proof of Lemma 3.2. To summarize, we know that

- $\hat{H}_P^{\varepsilon+1}(\Gamma) \cong H_N^{\varepsilon+1}(\Gamma) \cong 0$,
- every homogeneous cycle in $C_N^{\varepsilon}(\Gamma)$ can be completed to a cycle in $\hat{C}_P^{\varepsilon}(\Gamma)$ by adding terms with strictly lower polynomial degrees, and this correspondence gives rise to a well defined isomorphism $\hat{\phi}_n : H_N^{\varepsilon,n}(\Gamma) \to \mathcal{F}_x^n \hat{H}_P^{\varepsilon}(\Gamma)/\mathcal{F}_x^{n-1}\hat{H}_P^{\varepsilon}(\Gamma)$.

Clearly, $\pi_1(u_j)$ is a cycle in $\hat{C}_P^{\varepsilon}(\Gamma)$ obtained from $\pi_0(u_{j,n_j})$ by adding terms with polynomial degrees strictly less than $\deg \pi_0(u_{j,n_j}) = n_j$. Denote by $\pi^{(n)} : \mathcal{F}_x^n \hat{H}_P^{\varepsilon}(\Gamma) \to \mathcal{F}_x^n \hat{H}_P^{\varepsilon}(\Gamma)/\mathcal{F}_x^{n-1}\hat{H}_P^{\varepsilon}(\Gamma)$ the standard quotient map. Then $\pi^{(n_j)}(\pi_1([u_j])) = \hat{\phi}_{n_j}([\pi_0(u_{j,n_j})]) = \hat{\phi}_{n_j}(\pi_0([u_j])) \neq 0$. Thus, $\pi_1([u_j]) \notin \mathcal{F}_x^{n_j-1}\hat{H}_P^{\varepsilon}(\Gamma)$ and, therefore, $\deg_x \pi_1([u_j]) = n_j$.

It remains to show that $\sum_j c_j\pi_1([u_j]) \in \mathcal{F}_x^l \hat{H}_P(\Gamma)$ if and only if $c_j = 0$ whenever $n_j > l$. Recall that $\deg_x \pi_1([u_j]) = n_j$. If $c_j = 0$ whenever $n_j > l$, then we clearly have that $\sum_j c_j\pi_1([u_j]) \in \mathcal{F}_x^l \hat{H}_P(\Gamma)$. Now assume that there is at least one j such that $n_j > l$ and $c_j \neq 0$. Then $n := \max\{n_j \mid c_j \neq 0\} > l$.

Consider $[u] := \sum_j c_j a^{n-n_j}[u_j]$. $[u]$ is a homogeneous element of $H_P(\Gamma)$ of polynomial degree n, and $\pi_1([u]) = \sum_j c_j \pi_1([u_j])$. We have

$$\pi^{(n)}(\pi_1([u])) = \pi^{(n)}(\sum_{n_j=n} c_j \pi_1([u_j])) = \hat{\phi}_n(\sum_{n_j=n} c_j \pi_0([u_j])).$$

By Lemma 4.6, $\{\pi_0([u_j])\}$ is a basis for $H_N(\Gamma)$. By the definition of n, we know that $n_j = n$ and $c_j \neq 0$ for at least one j. Thus, $\sum_{n_j=n} c_j \pi_0([u_j]) \neq 0$. But $\hat{\phi}_n : H_N^{\varepsilon,n}(\Gamma) \to \mathcal{F}_x^n \hat{H}_P^{\varepsilon}(\Gamma)/\mathcal{F}_x^{n-1}\hat{H}_P^{\varepsilon}(\Gamma)$ is an isomorphism. So $\pi^{(n)}(\pi_1([u])) = \hat{\phi}_n(\sum_{n_j=n} c_j \pi_0([u_j])) \neq 0$. Thus $\sum_j c_j \pi_1([u_j]) = \pi_1([u]) \notin \mathcal{F}_x^l \hat{H}_P(\Gamma)(\subset \mathcal{F}_x^{n-1}\hat{H}_P(\Gamma))$. □

Remark 4.11. In conclusion of this subsection, we note that Lemmas 4.8 and 4.10 imply that any direct sum decomposition of $H_P(\Gamma)$ in the category of graded $\mathbb{C}[a]$-modules is also a direct sum decomposition in the category of filtered $\mathbb{C}$-spaces and induces a direct sum decomposition of $\hat{H}_P(\Gamma)$ in the category of filtered $\mathbb{C}$-spaces.

4.2. **Decomposition of $H_P(L)$.**

In this subsection, we give a proof of Lobb's decomposition theorem (Theorem 1.7.)

As before, a is a homogeneous variable of degree $2k$.

Lemma 4.12. [8] *Assume that* $(C^*, d^*) = \cdots \xrightarrow{d^{n-1}} C^n \xrightarrow{d^n} C^{n+1} \xrightarrow{d^{n+1}} \cdots$ *is a bounded chain complex of finitely generated graded-free* $\mathbb{C}[a]$*-module and its differential* d^* *preserves the grading of* C^**. Then, in the category of chain complexes of graded* $\mathbb{C}[a]$*-modules,* (C^*, d^*) *is a direct sum of chain complexes of the forms* $F_{i,s}$ *and* $T_{i,m,s}$ *given in* (1.5)–(1.6).

Proof. We prove the lemma by an induction on the total rank of C^*. If $\text{rank}C^* = 0$ or 1, then the lemma is trivially true. Assume $\text{rank}C^* = K \geq 2$ and the lemma is true if the rank of the chain complex is less than K. There is an n such that $C^n \neq 0$ and $C^j = 0 \ \forall \ j < n$. Consider the section $0 \to C^n \xrightarrow{d^n} C^{n+1} \xrightarrow{d^{n+1}} \cdots$ of C^*. Let $\{u_1, \ldots, u_p\}$ be a homogeneous basis for C^n with $\deg u_1 \leq \cdots \leq \deg u_p$, and $\{v_1, \ldots, v_q\}$ a homogeneous basis for C^{n+1} with $\deg v_1 \geq \cdots \geq \deg v_q$. For each $1 \leq j \leq p$, we have $d^n(u_j) = \sum_{i=1}^q f_{i,j} v_i$, where each $f_{i,j}$ is a monomial in a of degree $\deg f_{i,j} = \deg u_j - \deg v_i$. Note that $\deg f_{i,j}$ is increasing with respect to both i and j.

If $(f_{1,1}, \ldots, f_{q,1}) = 0$, then C^* has a direct sum component $0 \to \mathbb{C}[a] \cdot u_1 \to 0 \cong F_{n, \deg u_1}$. Thus, by induction hypothesis, the lemma is true for C^*.

If $(f_{1,1}, \ldots, f_{q,1}) \neq 0$, then there is an l such that $f_{l,1} \neq 0$ and $f_{i,1} = 0 \ \forall \ 1 \leq i < l$. Define a $q \times q$ matrix $\Xi = (\xi_{i,j})$ by

$$\xi_{i,j} = \begin{cases} 1 & \text{if } i = j, \\ \frac{f_{i,1}}{f_{l,1}} & \text{if } i > l \text{ and } j = l, \\ 0 & \text{otherwise.} \end{cases}$$

Then Ξ is invertible and $\Xi^{-1} = (\tilde{\xi}_{i,j})$ is given by

$$\tilde{\xi}_{i,j} = \begin{cases} 1 & \text{if } i = j, \\ -\frac{f_{i,1}}{f_{l,1}} & \text{if } i > l \text{ and } j = l, \\ 0 & \text{otherwise.} \end{cases}$$

Note that, for any $i > l$, $\frac{f_{i,1}}{f_{l,1}}$ is a monomial of a of degree $\deg f_{i,1} - \deg f_{l,1} = \deg v_l - \deg v_i$. We have

$$(d^n u_1, \ldots, d^n u_p) = (v_1, \ldots, v_q)(f_{i,j}) = (v_1, \ldots, v_q)\Xi\Xi^{-1}(f_{i,j}).$$

Let $\tilde{v}_j = \sum_{i=1}^q \xi_{i,j} v_i$ and $g_{i,j} = \sum_{\alpha=1}^q \xi_{i,\alpha} f_{\alpha,j}$. Then $\{\tilde{v}_1, \ldots, \tilde{v}_q\}$ is a homogeneous basis for C^{n+1} with $\deg \tilde{v}_i = \deg v_i$, $g_{i,j}$ is a monomial in a with $\deg g_{i,j} = \deg f_{i,j}$, and $(d^n u_1, \ldots, d^n u_p) = (\tilde{v}_1, \ldots, \tilde{v}_q)(g_{i,j})$.

Note that $g_{l,1} = f_{l,1} \neq 0$ and $f_{i,1} = 0 \ \forall \ i \neq l$. Next, define a $p \times p$ matrix $\Theta = (\theta_{i,j})$ by

$$\theta_{i,j} = \begin{cases} 1 & \text{if } i = j, \\ -\frac{g_{l,j}}{g_{l,1}} & \text{if } i = 1 \text{ and } j > 1, \\ 0 & \text{otherwise,} \end{cases}$$

where $\frac{g_{l,j}}{g_{l,1}}$ is a monomial of a of degree $\deg g_{l,j} - \deg g_{l,1} = \deg u_j - \deg u_1$. Let $\tilde{u}_j = \sum_{i=1}^{p} \theta_{i,j} u_i$ and $h_{i,j} = \sum_{\alpha=1}^{p} g_{i,\alpha}\theta_{\alpha,j}$. Then $\{\tilde{u}_1, \dots, \tilde{u}_p\}$ is a homogeneous basis for C^n with $\deg \tilde{u}_j = \deg u_j$, $h_{i,j}$ is a monomial in a with $\deg h_{i,j} = \deg g_{i,j} = \deg f_{i,j}$, and $(d^n\tilde{u}_1, \dots, d^n\tilde{u}_p) = (\tilde{v}_1, \dots, \tilde{v}_q)(h_{i,j})$.

Note that $h_{l,1} = g_{l,1} = f_{l,1} \neq 0$, $h_{i,1} = 0 \ \forall\ i \neq l$ and $h_{l,j} = 0 \ \forall\ j \neq 1$. Thus, C^* has a direct sum component

$$0 \to \mathbb{C}[a] \cdot \tilde{u}_1 \xrightarrow{h_{l,1}} \mathbb{C}[a] \cdot \tilde{v}_l \to 0 \cong T_{n+1, \frac{\deg \tilde{u}_1 - \deg \tilde{v}_l}{2k}, \deg \tilde{v}_l}.$$

By the induction hypothesis, the lemma is true for C^*. □

The existence of the decomposition in Theorem 1.7 follows from Lemma 4.12. To prove the uniqueness of this decomposition, we need the following lemma, which is a slight refinement of the standard invariance theorem for modules over a principal ideal domain.

Lemma 4.13. *Suppose that* $\{(m_1, s_1), \dots, (m_p, s_p)\}$ *and* $\{(n_1, t_1), \dots, (n_q, t_q)\}$ *are two sequences in* $\mathbb{Z}_{>0} \times \mathbb{Z}$ *satisfying:*

- $m_1 \leq \cdots \leq m_p$, $n_1 \leq \cdots \leq n_q$.
- *If* $i < j$ *and* $m_i = m_j$, *then* $s_i \leq s_j$.
- *If* $i < j$ *and* $n_i = n_j$, *then* $t_i \leq t_j$.
- *As graded* $\mathbb{C}[a]$*-modules,*

$$\bigoplus_{i=1}^{p} (\mathbb{C}[a]/(a^{m_i}))\{s_i\} \cong \bigoplus_{j=1}^{p} (\mathbb{C}[a]/(a^{n_j}))\{t_j\}.$$

Then $p = q$ *and* $m_i = n_i$, $s_i = t_i$ *for every* $1 \leq i \leq p$.

Proof. We adapt the proof of the invariance theorem in [3, Section 3.9] to prove Lemma 4.13. The only change is that, instead of counting dimensions, we count graded dimensions. Recall that, for a finite dimensional graded $\mathbb{C}$-space $V = \bigoplus_i V_i$, where V_i is the homogeneous component of V of degree i, the graded dimension of V is $\mathrm{gdim}_{\mathbb{C}} V := \sum_i \beta^i \dim_{\mathbb{C}} V_i$, where β is a homogeneous variable of degree 1.

Let

$$M := \bigoplus_{i=1}^{p} (\mathbb{C}[a]/(a^{m_i}))\{s_i\} \cong \bigoplus_{j=1}^{p} (\mathbb{C}[a]/(a^{n_j}))\{t_j\}.$$

Denote by z_i the multiplicative unit 1 in $(\mathbb{C}[a]/(a^{m_i}))\{s_i\}$, which is a homogeneous element of M of degree s_i. For any $l \geq 0$, define $M^{(l)} := a^l M / a^{l+1} M$. Then $M^{(l)}$ is a finite dimensional graded $\mathbb{C}$-space. If $l \geq m_p$, then $M^{(l)} = 0$ and $\mathrm{gdim}_{\mathbb{C}} M^{(l)} = 0$. If $0 \leq l < m_p$, then there is a unique j such that $m_1 \leq \cdots \leq m_j \leq l < m_{j+1} \leq \cdots \leq m_p$. One can see that $a^l M = \bigoplus_{i=j+1}^{p} \mathbb{C}[a] a^l z_i$ and $\{a^l z_{j+1} + a^{l+1}M, \dots, a^l z_p + a^{l+1}M\}$ is a homogeneous basis for the $\mathbb{C}$-space $M^{(l)}$. So $\mathrm{gdim}_{\mathbb{C}} M^{(l)} = \beta^{2kl} \sum_{i=j+1}^{p} \beta^{s_i}$. For any integer c, define $S_{l,c} = \{i \mid 1 \leq i \leq p,\ s_i = c,\ m_i > l\}$. We observe that the coefficient of β^{c+2kl} in $\mathrm{gdim}_{\mathbb{C}} M^{(l)}$ is equal to the cardinality of $S_{l,c}$. Similarly, defining $S'_{l,c} = \{i \mid 1 \leq i \leq q,\ t_i = c,\ n_i > l\}$, we have that the coefficient of β^{c+2kl} in $\mathrm{gdim}_{\mathbb{C}} M^{(l)}$ is equal to the cardinality of $S'_{l,c}$. Thus, for any $(l, c) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}$, the cardinalities of $S_{l,c}$ and $S'_{l,c}$ are equal. The lemma follows from this. □

It is now very easy to prove Theorem 1.7 and Corollary 1.8.

Proof of Theorem 1.7 and Corollary 1.8. Fix a diagram D of L and a marking of D. Let $x_1, \dots, x_m$ be the variables assigned to the marked points of Γ and $R = \mathbb{C}[x_1, \dots, x_m, a]$. By Corollary 3.3, $H(C_P(D), d_{mf})$ with its polynomial grading is a finitely generated graded-free $\mathbb{C}[a]$-module. By the definition of d_χ, we know that it preserves the polynomial grading.

According to Lemma 4.12, in the category of chain complexes of graded $\mathbb{C}[a]$-modules, $(H(C_P(D), d_{mf}), d_\chi)$ decomposes into a direct sum of chain complexes of the forms $F_{i,s}$ and $T_{i,m,s}$ given in (1.5)–(1.6). Note that each factor of $F_{i,s}$ in this decomposition contributes a direct sum component $\mathbb{C}[a]\|i\|\{s\}$ to $H_P(L)$ and each factor of $T_{i,m,s}$ contributes a direct sum component $(\mathbb{C}[a]/(a^m))\|i\|\{s\}$ to $H_P(L)$. To prove the existence of decomposition (1.7), it remains to determine the free part of $H_P(L)$. By Lemma 4.10, the above decomposition of $(H(C_P(D), d_{mf}), d_\chi)$ induces a decomposition of $(H(\hat{C}_P(D), d_{mf}), d_\chi)$ in the category of filtered chain complexes of $\mathbb{C}$-spaces. Each factor of $F_{i,s}$ (resp. $T_{i,m,s}$) in the decomposition of $(H(C_P(D), d_{mf}), d_\chi)$

corresponds to a factor of $\hat{F}_{i,s}$ (resp. $\hat{T}_{i,m,s}$) in the decomposition of $(H(\hat{C}_P(D), d_{mf}), d_\chi)$, and the grading of $F_{i,s}$ (resp. $T_{i,m,s}$) induces the filtration on the corresponding $\hat{F}_{i,s}$ (resp. $\hat{T}_{i,m,s}$). The homology of $\hat{T}_{i,m,s}$ vanishes and, as a filtered $\mathbb{C}$-space, the homology of $\hat{F}_{i,s}$ is $\mathbb{C}\|i\|\{s\} \cong (\mathbb{C}[a]/(a-1))\|i\|\{s\}$. One can see from this that, as a graded $\mathbb{C}[a]$-module, the free part of $H_P(L)$ is isomorphic to $\hat{\mathcal{H}}_P(L) \otimes_{\mathbb{C}} \mathbb{C}[a]$. Thus, we have proved the existence of decomposition (1.7).

The uniqueness of decomposition (1.7) follows from Lemma 4.13. This completes the proof of Theorem 1.7.

To prove Corollary 1.8, note that, by Lemma 4.6, $(H(C_N(D), d_{mf}), d_\chi) \cong H(C_P(D), d_{mf})/aH(C_P(D), d_{mf})$ as chain complexes of graded $\mathbb{C}$-spaces. Moreover, as chain complexes of graded $\mathbb{C}$-spaces, $F_{i,s}/aF_{i,s} \cong 0 \to \mathbb{C}\|i\| \to 0$ and $T_{i,m,s}/aT_{i,m,s} \cong 0 \to \mathbb{C}\|i-1\|\{s+2km\} \xrightarrow{0} \mathbb{C}\|i\|\{s\} \to 0$. So Corollary 1.8 also follows from the decomposition of the chain complex $(H(C_P(D), d_{mf}), d_\chi)$. $\square$

4.3. **Decompositions of $E_r(L)$ and $\hat{E}_r(L)$.** In this subsection, we prove the decompositions of $E_r(L)$ and $\hat{E}_r(L)$.

First, we explicitly compute the spectral sequences of the filtered chain complexes $F_{i,s}$, $T_{i,m,s}$, $\hat{F}_{i,s}$ and $\hat{T}_{i,m,s}$. We use the notations given in [9, Section 2.2]. Since $\mathcal{F}_x$ is an increasing filtration and the notations in [9, Section 2.2] are for a decreasing filtration, we need to adjust their definitions accordingly.

Let $(C^*, d, \mathcal{F})$ be a filtered chain complex such that d raises the homological grading by 1 and $\mathcal{F}$ is increasing. Set

$$Z_r^{p,q} = \mathcal{F}^p C^{p+q} \cap d^{-1}(\mathcal{F}^{p-r} C^{p+q+1}), \tag{4.1}$$

$$B_r^{p,q} = \mathcal{F}^p C^{p+q} \cap d(\mathcal{F}^{p+r} C^{p+q-1}), \tag{4.2}$$

$$E_r^{p,q} = Z_r^{p,q}/(Z_{r-1}^{p-1,q+1} + B_{r-1}^{p,q}). \tag{4.3}$$

Then $\{E_r^{p,q}\}$ is the spectral sequence of $(C^*, d, \mathcal{F})$.

Proof of Lemma 1.10. We only compute $E_r(F_{i,s})$ and $E_r(T_{i,m,s})$. The computation of $E_r(\hat{F}_{i,s})$ and $E_r(\hat{T}_{i,m,s})$ is very similar and left to the reader.

Recall that the differential map of $F_{i,s}$ is 0. So $B_r^{p,q}(F_{i,s}) = 0$ and

$$Z_r^{p,q}(F_{i,s}) = \mathcal{F}^p F_{i,s}^{p+q} = \begin{cases} \mathbb{C}[a]\{s\} & \text{if } p \geq s,\ q = i-p, \\ 0 & \text{otherwise.} \end{cases}$$

So

$$E_r^{p,q}(F_{i,s}) = \mathcal{F}^p F_{i,s}^{p+q}/\mathcal{F}^{p-1} F_{i,s}^{p+q} \cong \begin{cases} \mathbb{C}[a]\{s\} & \text{if } p = s,\ q = i-s, \\ 0 & \text{otherwise.} \end{cases}$$

The observations about $\{E_r(F_{i,s})\}$ in Lemma 1.10 follow from this.

Recall that the filtered chain complex $T_{i,m,s}$ is given by

$$\mathcal{F}_x^p T_{i,m,s} = \begin{cases} 0 \to \mathbb{C}[a]\|i-1\|\{s+2km\} \xrightarrow{a^m} \mathbb{C}[a]\|i\|\{s\} \to 0 & \text{if } p \geq s+2km, \\ 0 \to \mathbb{C}[a]\|i\|\{s\} \to 0 & \text{if } s \leq p < s+2km, \\ 0 & \text{if } p < s. \end{cases}$$

Note that $Z_r^{p,q}(T_{i,m,s}) = 0$ unless $q = i-1-p$ or $i-p$. So $E_r^{p,q}(T_{i,m,s}) = 0$ unless $q = i-1-p$ or $i-p$.

We compute $E_r^{p,i-1-p}(T_{i,m,s})$ first. Note that

$$\mathcal{F}_x^p T_{i,m,s}^{i-1} = \begin{cases} \mathbb{C}[a]\{s+2km\} & \text{if } p \geq s+2km, \\ 0 & \text{if } p < s+2km, \end{cases}$$

$$d^{-1}(\mathcal{F}_x^{p-r} T_{i,m,s}^{i}) = \begin{cases} \mathbb{C}[a]\{s+2km\} & \text{if } p-r \geq s, \\ 0 & \text{if } p-r < s. \end{cases}$$

Thus,

$$Z_r^{p,i-1-p}(T_{i,m,s}) = \mathcal{F}_x^p T_{i,m,s}^{i-1} \cap d^{-1}(\mathcal{F}_x^{p-r} T_{i,m,s}^{i}) = \begin{cases} \mathbb{C}[a]\{s+2km\} & \text{if } p \geq s+2km \text{ and } p \geq s+r \\ 0 & \text{otherwise.} \end{cases}$$

Also, $B_r^{p,i-1-p}(T_{i,m,s}) = \mathcal{F}_x^p T_{i,m,s}^{i-1} \cap d(\mathcal{F}_x^{p+r} T_{i,m,s}^{i-2}) = 0$. Therefore,

$$E_r^{p,i-1-p}(T_{i,m,s}) = Z_r^{p,i-1-p}(T_{i,m,s})/Z_r^{p-1,i-p}(T_{i,m,s}) \cong \begin{cases} \mathbb{C}[a]\{s+2km\} & \text{if } p = s+2km \geq s+r \\ 0 & \text{otherwise.} \end{cases}$$

Next, we compute $E_r^{p,i-p}(T_{i,m,s})$. Note that

$$Z_r^{p,i-p}(T_{i,m,s}) = \mathcal{F}_x^p T_{i,m,s}^i \cap d^{-1}(\mathcal{F}_x^{p-r} T_{i,m,s}^{i+1}) = \mathcal{F}_x^p T_{i,m,s}^i = \begin{cases} \mathbb{C}[a]\{s\} & \text{if } p \geq s, \\ 0 & \text{if } p < s, \end{cases}$$

and

$$d(\mathcal{F}_x^{p+r} T_{i,m,s}^{i-1}) = \begin{cases} a^m \mathbb{C}[a]\{s\} & \text{if } p+r \geq s+2km, \\ 0 & \text{if } p+r < s+2km. \end{cases}$$

So

$$B_r^{p,i-p}(T_{i,m,s}) = \begin{cases} a^m \mathbb{C}[a]\{s\} & \text{if } p \geq s \text{ and } p+r \geq s+2km, \\ 0 & \text{if } p+r < s+2km. \end{cases}$$

Recall that $E_r^{p,i-p}(T_{i,m,s}) = Z_r^{p,i-p}(T_{i,m,s})/(Z_r^{p-1,i-p+1}(T_{i,m,s}) + B_r^{p,i-p}(T_{i,m,s}))$. Putting these together, we get

$$E_r^{p,i-p}(T_{i,m,s}) \cong \begin{cases} (\mathbb{C}[a]/(a^m))\{s\} & \text{if } p = s \text{ and } r \geq 2km+1, \\ \mathbb{C}[a]\{s\} & \text{if } p = s \text{ and } r \leq 2km, \\ 0 & \text{otherwise.} \end{cases}$$

This completes the computation of $\{E_r(T_{i,m,s})\}$. Note that $\{E_r(T_{i,m,s})\}$ collapses exactly at its E_{2km+1}-page. □

Finally, we prove Theorem 1.11.

Proof of Theorem 1.11. Fix a diagram D of L and a marking of D. Let $x_1, \dots, x_m$ be the variables assigned to the marked points of Γ and $R = \mathbb{C}[x_1, \dots, x_m, a]$. By Corollary 3.3, $H(C_P(D), d_{mf})$ with its polynomial grading is a finitely generated graded-free $\mathbb{C}[a]$-module. By the definition of d_χ, we know it preserves the polynomial grading.

According to Lemma 4.12, in the category of chain complexes of graded $\mathbb{C}[a]$-modules, $(H(C_P(D), d_{mf}), d_\chi)$ decomposes into a direct sum of chain complexes of the forms $F_{i,s}$ and $T_{i,m,s}$ given in (1.5)–(1.6). By Lemma 4.8, this is also a decomposition in the category of filtered chain complexes, in which the filtrations on $F_{i,s}$ and $T_{i,m,s}$ are given by (1.9)–(1.10). Thus, the spectral sequence of $(H(C_P(D), d_{mf}), d_\chi)$ is the direct sum of the spectral sequences of its components in this decomposition. Decomposition (1.18) in Theorem 1.11 then follows from this and Lemma 1.10.

By Lemma 4.10, the above decomposition of $(H(C_P(D), d_{mf}), d_\chi)$ induces a decomposition of $(H(\hat{C}_P(D), d_{mf}), d_\chi)$ in the category of filtered chain complexes into a direct sum of chain complexes of the forms $\hat{F}_{i,s}$ and $\hat{T}_{i,m,s}$ given in (1.11)–(1.12). The spectral sequence of $(H(\hat{C}_P(D), d_{mf}), d_\chi)$ is the direct sum of the spectral sequences of its components in this decomposition. Decomposition (1.17) in Theorem 1.11 then follows from this and Lemma 1.10. □

REFERENCES

[1] A. Elliott, *Graphical methods establishing nontriviality of state cycle Khovanov homology classes,* arXiv:0907.0396.

[2] B. Gornik, *Note on Khovanov link cohomology,* arXiv:math.QA/0402266.

[3] N. Jacobson, *Basic algebra. I,* Second edition. W. H. Freeman and Company, New York, 1985. xviii+499 pp. ISBN: 0-7167-1480-9 00A05 (12-01)

[4] M. Khovanov, L. Rozansky, *Matrix factorizations and link homology,* Fund. Math. **199** (2008), no. 1, 1–91.

[5] D. Krasner, *Equivariant sl(n)-link homology,* Algebr. Geom. Topol. 10 (2010), no. 1, 1-32.

[6] P. Kronheimer, T. Mrowka, *Khovanov homology is an unknot-detector,* Publ. Math. Inst. Hautes tudes Sci. No. 113 (2011), 97-208.

[7] E. Lee, *An endomorphism of the Khovanov invariant,* Adv. Math. **197** (2005), no. 2, 554–586.

[8] A. Lobb, 2*-strand twisting and knots with identical quantum knot homologies,* arXiv:1103.1412v3.

[9] J. McCleary, *A user's guide to spectral sequences.,* Second edition. Cambridge Studies in Advanced Mathematics, 58. Cambridge University Press, Cambridge, 2001. xvi+561 pp. ISBN: 0-521-56759-9

[10] J. Rasmussen, *Khovanov homology and the slice genus,* Invent. Math. **182** (2010), no. 2, 419-447.
[11] H. Wu, *On the quantum filtration of the Khovanov-Rozansky cohomology,* Adv. Math. **221** (2009), no. 1, 54–139.
[12] H. Wu, *A colored* $\mathfrak{sl}(N)$*-homology for links in* S^3*,* arXiv:0907.0695v5.
[13] H. Wu, *Equivariant colored* $\mathfrak{sl}(N)$*-homology for links,* J. Knot Theory Ramifications **21**, No. 2 (2012), 1250012 (104 pages.)
[14] H. Wu, *Generic deformations of the colored* $\mathfrak{sl}(N)$*-homology for links,* Algebr. Geom. Topol. **11** (2011) 2037–2106.
[15] H. Wu, *Simplified Khovanov-Rozansky chain complexes for open* 2*-braids,* Topology Appl. **159**, No. 14 (2012), 3190-3203.

DEPARTMENT OF MATHEMATICS, THE GEORGE WASHINGTON UNIVERSITY, MONROE HALL, ROOM 240, 2115 G STREET, NW, WASHINGTON DC 20052, USA. TELEPHONE: 1-202-994-0653, FAX: 1-202-994-6760
E-mail address: `haowu@gwu.edu`